

MRN: \_\_\_\_\_

**Palmetto Infusion Services  
Standard Plan of Treatment for Lumizyme**

(Re)Certification Period From \_\_\_\_\_ to \_\_\_\_\_

**Note:** We may require MD office notes and may require a letter of Medical Necessity (depending on diagnosis), to be able to verify eligibility and payment for this treatment through patients Medicare and/or other insurance plan.

**Patient's Name** \_\_\_\_\_ **HT:** \_\_\_\_\_ **Weight:** \_\_\_\_\_ **Allergies** \_\_\_\_\_

Primary Diagnosis: Pompe's Disease Secondary Diagnosis: \_\_\_\_\_

**ORDERS:**

Obtain weight each visit. **Vital Signs every 30 minutes beginning with start of infusion and with each rate change, at completion of infusion and after 1 hour wait.**

Instruct patient/caregiver on medications, signs/symptoms of adverse reaction. Assess patient response to therapy.

Utilize existing central line for administration, or initiate a peripheral IV with each infusion, prn Normal Saline Flush 5 ml before and 20 ml flush thru tubing after infusion followed by Heparin Lock 1-5ml 100 units/ml as needed per line type.

Pump, tubing, and supplies needed to complete prescribed therapy.

**If adverse drug reaction, implement the Standing Adverse Reaction protocol.**

**Drug:**

Lumizyme: \_\_\_\_\_ Mg in \_\_\_\_\_ ml 0.9% Normal Saline, every \_\_\_\_\_ Wks

Rate of infusion as follows:

Step 1 \_\_\_\_\_/hr x 30 mins                      Step 2 \_\_\_\_\_/hr x 30 mins

Step 3 \_\_\_\_\_/hr x 30 mins                      Step 4 \_\_\_\_\_/hr x 30 mins

**Lab Orders:**

IGG level every 3 months or: \_\_\_\_\_ (specify frequency)

Pharmacist to perform clinical drug monitoring.

(No Stamped Signatures please)

Physicians Signature: \_\_\_\_\_ Date: \_\_\_\_\_

(Substitution Permitted)

**PLEASE FAX DEMOGRAPHICS AND INSURANCE INFORMATION TO:  
TOLL FREE FAX (866) 872-8920**

\_\_\_\_\_