



MRN: _____

DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Guidelines for Prescribing HyQvia®
(Required documentation with all initial referrals)

Patient Name: _____

Referral Date: _____

- _____ Include signed and completed **Plan of Treatment**. (MD must complete sections 1-6)
(Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab)
- _____ Include patient demographic information and insurance information. (Copy of insurance cards if available)
- _____ **Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include IG levels, other lab results and/or tests to support diagnosis.**
 - HyQvia® is an immune globulin with a Recombinant Human Hyaluronidase indicated for the treatment of Primary Immunodeficiency (PI) in adults. This includes, but is not limited to, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, congenital agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies^{1, 2}.
- _____ If patient is currently on IVIG therapy, then please specify therapy and last dose: _____
 - For patients previously on another IgG treatment, it is recommended to administer the first dose approximately one week after the last infusion of their previous treatment.
- _____ Other as requested: _____

Pre-Screening:
_____ **IG levels**

**** Warnings/Precautions: IgA-deficient patients with anti-IgA antibodies** are at greater risk of severe hypersensitivity and anaphylactic reactions. • **Thrombosis** may occur with immune globulin products, including HYQVIA. Risk factors may include advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling vascular catheters, hyper viscosity and cardiovascular risk factors. • **Antibodies to PH20** (Recombinant Human Hyaluronidase) can develop. The potential exists for such antibodies to cross-react with endogenous PH20 which is known to be expressed in the adult male testes, epididymis, and sperm. It is unknown whether these antibodies may interfere with fertilization in humans. • **Aseptic Meningitis Syndrome (AMS)** may occur. Discontinue treatment if AMS symptoms appear. • **Acute intravascular hemolysis** may occur. Monitor for clinical signs and symptoms of hemolysis and hemolytic anemia. • Infusion into or around an infected area can spread a localized infection. • **Monitor for pulmonary adverse reactions** (transfusion-related acute lung injury [TRALI]). • **May carry a risk of transmitting infectious agents**, e.g., viruses, the variant Creutzfeldt-Jakob disease (vCJD) agent and, theoretically, the CreutzfeldtJakob disease (CJD) agent. • **Acute renal dysfunction/failure** has been reported in association with Immune Globulin Infusion 10% (Human) administered intravenously. Ensure that patients are not volume depleted prior to the initiation of infusion of HYQVIA. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.