



Phone: 1-800-809-1265 Fax: 1-866-872-8920

MRN: _____

DOB: _____

STANDARD CYCLOPHOSPHAMIDE PLAN OF TREATMENT

NOTE: Patient may be ineligible to receive cyclophosphamide if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, unable to adequately hydrate, or planned/recent surgery.

1. Patient Name: _____ Height (inches): _____ Weight (lbs): _____

2. Allergies: _____

3. Diagnosis: Primary ICD-10 Code: _____ **Diagnosis description:** _____

Secondary ICD-10 Code: _____ **Diagnosis description:** _____

4. Pre-medications: None **OR Administered 30 minutes prior to infusion as selected:**

<p>Acetaminophen:</p> <p><input type="checkbox"/> 650 mg PO</p> <p><input type="checkbox"/> 500 mg PO</p> <p><input type="checkbox"/> 325 mg PO</p>	<p>Ondansetron: <input type="checkbox"/> 4 mg PO, <input type="checkbox"/> 4 mg IV, <input type="checkbox"/> 8 mg PO, or <input type="checkbox"/> 8 mg IV. May repeat x _____</p> <p><input type="checkbox"/> 10 mg/50 ml Sodium Chloride IV over 15-30 minutes</p> <p><input type="checkbox"/> Dexamethasone 10 mg IV over 15-30 mins or other _____ mg IV</p> <p>Mesna _____ mg over _____ minutes IV prior to infusion</p>
---	---

Pre-medicate with other: _____

CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion

Check for blood return prior to start of Cyclophosphamide and utilize Hazardous Drug Precautions.

- 5. Orders:**
- Cyclophosphamide _____ mg in _____ ml of Sodium Chloride 0.9% IV to infuse over _____ minutes
 - Pre- hydration: Infuse _____ ml Sodium Chloride 0.9% IV over _____ hours before Cyclophosphamide
 - Post- hydration: Infuse _____ ml Sodium Chloride 0.9% IV over _____ hours after Cyclophosphamide
 - Encourage oral hydration of _____ after infusion

6. Frequency: Orders to be completed every _____ week (s) x _____ month(s)

Special Orders: _____

Clinical lab monitoring required: CBC results required 7-10 days after each infusion. Next treatment will be held for WBC level less than 2.5 K/uL. The referring MD office will be notified, and treatment will not be resumed until MD clearance received.

If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES

7. Physician's Signature: _____ / _____ Date: _____
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: _____ NPI: _____

8. Fax updated supporting clinical MD notes with each order renewal or change in orders
Infusion order forms and Adverse Drug Reaction Guidelines are available at www.palmettoinfusion.com

