



MRN: _____

DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

STANDARD FASENRA™ (benralizumab) PLAN OF TREATMENT

NOTE: Patient may be ineligible to receive FASENRA™ (benralizumab) if patient has signs/symptoms of a parasitic infection, is currently being treated for a parasitic infection, or is having an acute bronchospasm and/or asthma attack.

1. Patient Name: _____ Height (inches): _____ Weight (lbs): _____

2. Allergies: _____

3. Diagnosis:

- J45.51 Severe persistent asthma with (acute) exacerbation J45.50 Severe persistent asthma, uncomplicated
- Other ICD-10 Code: _____ Diagnosis description: _____

CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion

4. Dose/Frequency: (Administer as subcutaneous injection only to upper arm, thigh, or abdomen)

Induction dose:

FASENRA™ (benralizumab) 30 mg subcutaneous injection every 4 weeks for the first (3) doses given at week 0, week 4, week 8 and then followed by once every 8 weeks thereafter.

Maintenance dose as follows:

FASENRA™ (benralizumab) 30 mg subcutaneous injection every 8 weeks.

Extended post treatment monitoring for any patient new to therapy: monitor patient for one (1) hour after first injection, for 30- minutes after second injection, and then 15-minutes after each injection thereafter.

Special orders: _____

Patient to continue Fasenra unless:

- Previous allergic reaction
- Parasitic infection
- Pregnancy
- Other: _____

If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES

5. Physician's Signature: _____ / _____ Date: _____
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: _____ NPI: _____

6. Fax updated supporting clinical MD notes with each order renewal or change in orders
Infusion order forms and Adverse Drug Reaction Guidelines are available at www.palmettoinfusion.com



MRN: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: _____

Guidelines for Prescribing FASENRA™ (benralizumab)

(Required documentation with all initial referrals)

Patient Name: _____ Referral Date: _____

- Include signed and completed **Plan of Treatment**. (MD must complete sections 1-6)
(Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab)
- Include patient demographic information and insurance information. (Copy of insurance cards if available)
- Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or Pulmonary Function Tests to support diagnosis.**
 - FASENRA™ (benralizumab) is an interleukin-5 receptor alpha-directed cytolytic monoclonal antibody (IgG1, kappa) indicated for the **add-on maintenance treatment** of patients with severe asthma aged 12 years and older, and with an eosinophilic phenotype.
- If patient is switching biologic therapies such as Xolair®, Cinqair®, or Nucala®, then MD must specify wash-out period prior to starting **FASENRA™** as specified of _____ weeks.
Last known therapy: _____ and last known date received: _____.
- Other as requested: _____

Pre-Screening:

- Blood Eosinophil Level

**** Warnings/Precautions: Hypersensitivity reactions:** hypersensitivity reactions (e.g., anaphylaxis, angioedema, urticaria, rash) have occurred after administration of FASENRA™. Discontinue in the event of a hypersensitivity reaction. The most common adverse reactions (incidence greater than or equal to 5%) include headache and pharyngitis. • **Reduction in Corticosteroid Dosage:** Do not discontinue systemic or inhaled corticosteroids abruptly upon initiation of therapy. Decrease corticosteroids gradually, if appropriate. • **Parasitic (Helminth) Infection:** Treat patients with pre-existing helminth infections before therapy with FASENRA™. If patients become infected while receiving FASENRA™ and do not respond to anti-helminth treatment, discontinue FASENRA™ until the parasitic infection resolves. Pregnancy/Breastfeeding: Discuss Pregnancy or breastfeeding plans/risks prior to start of therapy. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.