



MRN: \_\_\_\_\_

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: \_\_\_\_\_

### STANDARD Orenzia® (abatacept) PLAN OF TREATMENT FOR RHEUMATOLOGY

**NOTE:** Patient *may be ineligible* to receive abatacept if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening diagnosis of COPD or respiratory status, and/or surgery.

1. Patient Name: \_\_\_\_\_ Height (inches): \_\_\_\_\_ Weight (lbs): \_\_\_\_\_

2. Allergies: \_\_\_\_\_

3. **Diagnosis:** \* Please complete the 2<sup>nd</sup> and 3<sup>rd</sup> digits to complete the ICD-10 code for billing

M05. \_\_\_\_\_ Rheumatoid Arthritis with Rheumatoid factor    M06. \_\_\_\_\_ Rheumatoid Arthritis without Rheumatoid factor

Other ICD-10 Code: \_\_\_\_\_ Diagnosis description: \_\_\_\_\_

4. **Pre-medications:**  None *OR* Administered 30 minutes prior to infusion as selected:

**\*Product information does not suggest premedication and suggest caution with use of Diphenhydramine due to 30 minute infusion time and safety risks with driving.**

<p>a) Acetaminophen:</p> <p><input type="checkbox"/> 650mgs PO</p> <p><input type="checkbox"/> 500mgs PO</p> <p><input type="checkbox"/> 325mgs PO</p>	<p>b) Diphenhydramine: <input type="checkbox"/> 25 mgs PO, <input type="checkbox"/> 50mgs PO, <input type="checkbox"/> 25 mgs IVP, <input type="checkbox"/> 50mgs IVP or</p> <p>c) Alternate oral antihistamine to diphenhydramine:</p> <p><input type="checkbox"/> Cetirizine 10 mg, <input type="checkbox"/> Loratadine 10 mg, Fexofenadine <input type="checkbox"/> 60mgs or <input type="checkbox"/> 180mgs</p> <p>d) Other: Methylprednisolone <input type="checkbox"/> 40mgs IVP <input type="checkbox"/> 125mgs IVP or other _____mgs IVP</p> <p>Famotidine: <input type="checkbox"/> 20mgs PO, <input type="checkbox"/> 40mgs PO, <input type="checkbox"/> 20mgs IVP, <input type="checkbox"/> 40mgs IVP</p>
--	--

e) Pre-medicate with other: \_\_\_\_\_

CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion

**Orders:** Obtain weight each visit. **Vital signs at baseline and after completion, then may discharge when infusion is complete.** Instruct patient/caregiver on medications and signs/symptoms of adverse reaction. Assess patient for response to therapy. Utilize existing central line for administration or initiate a peripheral IV with each infusion as needed. Sodium Chloride 0.9% flush 3-10 ml before, after, and as needed during the infusion. Follow infusion with Heparin 100 units/ml 1 – 5 ml per line type or to peripheral IV as required for multiple day treatments. Pump, tubing, 0.22 micron filter, and supplies needed to include silicone-free syringe for reconstitution to complete prescribed therapy. Pharmacist to perform clinical drug monitoring. **If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES.**

5. **Dose:** Orenzia® (abatacept) dosage per 100 ml Sodium Chloride 0.9% IV to infuse over at least 30 minutes.

Orenzia® (abatacept) dosage will be based on the following guidelines provided by Bristol-Myers Squibb

Body Weight of Patient	Dose	Number of 250mg (abatacept) Vials
< 60 kg	500 mg	2
60 to 100 kg	750 mg	3
>100 kg	1000 mg	4

6. **Frequency:** \_\_\_\_\_ Induction orders to be completed at 0 week, 2 week, and 4 weeks, and then every 4 weeks thereafter

\_\_\_\_\_ Orders every 4 weeks (maintenance).

\_\_\_\_\_ Special Orders: \_\_\_\_\_

Lab orders with infusions: \_\_\_\_\_

7. **Physician's Signature:** \_\_\_\_\_ / \_\_\_\_\_ Date: \_\_\_\_\_

No Stamp Signatures

(Dispense as written)

(Substitution permitted)

Printed Physician's Name with Credentials: \_\_\_\_\_

**8. Fax updated supporting clinical MD notes with each order renewal or change in orders**

*Infusion order forms available at [www.palmettoinfusion.com](http://www.palmettoinfusion.com)*



MRN: \_\_\_\_\_

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: \_\_\_\_\_

## Guidelines for Prescribing Orenzia® (abatacept) for Rheumatology

(Required documentation with all initial referrals)

Patient Name: \_\_\_\_\_

Referral Date: \_\_\_\_\_

\_\_\_ Include signed and completed **Plan of Treatment**. (MD must complete sections 1-8)  
 (Infusion order forms & Standard Adverse Reactions orders are available at [www.palmettoinfusion.com](http://www.palmettoinfusion.com) under Agency/MD tab)

\_\_\_ Include patient demographic information and insurance information. (Copy of insurance cards if available)

\_\_\_ **Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.**

- Orenzia® is indicated for moderately to severely active RA in adults. ORENCIA may be used as monotherapy or concomitantly with DMARDs other than TNF antagonists.

\_\_\_ If patient is switching biological therapies, then MD must specify wash-out period prior to starting Orenzia® as specified of \_\_\_\_\_ weeks. Last known biological therapy: \_\_\_\_\_ and last date received: \_\_\_\_\_. (Include copy of last Orenzia® infusion record if available and currently on therapy)

\_\_\_ Other as requested: \_\_\_\_\_

**Pre-Screening:** (TB and Hepatitis screening results must be available prior to start of therapy and within last 12 months.)

\_\_\_ **Required TB screening results: PPD or QuantiFERON Gold Test.**  
 (\* If screening results are positive or indeterminate, then a negative CXR result is required.)

\_\_\_ **Required Hepatitis screening to include: Hepatitis B Surface Antigen results.**

\*\* Warnings/Precautions: Patient should not have an active ongoing infection, signs or symptoms of malignancy, or invasive fungal infection. **COPD patients** may develop more frequent respiratory adverse events. Orenzia® in patients with RA and COPD should be undertaken with caution and such patients should be monitored for worsening of their respiratory status. **Blood Glucose Monitoring:** Orenzia® contains maltose and can interfere with the readings of blood glucose monitors that use test strips with (GDH-PQQ), resulting in falsely elevated blood glucose readings on the day of infusion. **Hepatitis B Reactivation:** Monitor HBV carriers during and several months after therapy. If reactivation occurs, stop Orenzia® and begin anti-viral therapy. Evaluation of immunizations should be completed prior to and **live vaccines** should not be given before or concurrently with Orenzia®. See full prescribing information.

**Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.**

**Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.**