



Phone: 1-800-809-1265 Fax: 1-866-872-8920

MRN: \_\_\_\_\_

DOB: \_\_\_\_\_

**STANDARD RENFLEXIS (infliximab-abda) PLAN OF TREATMENT FOR GASTROENTEROLOGY**

**NOTE:** Patient *may be ineligible* to receive (infliximab-abda) if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

**1. Patient Name:** \_\_\_\_\_ Height (inches): \_\_\_\_\_ Weight (lbs): \_\_\_\_\_

**2. Allergies:** \_\_\_\_\_

**3. Diagnosis:** \* Please complete the 2<sup>nd</sup> and 3<sup>rd</sup> digits to complete the ICD-10 code for billing

- K50.0** \_\_\_\_\_ Crohn's Disease (small intestine)
- K50.1** \_\_\_\_\_ Crohn's Disease (large intestine)
- K50.8** \_\_\_\_\_ Crohn's Disease (small & large intestine)
- K50.9** \_\_\_\_\_ Crohn's Disease, Unspecified
- K51.8** \_\_\_\_\_ Other Ulcerative (chronic) Colitis
- K51.5** \_\_\_\_\_ Left sided Ulcerative (chronic) Colitis
- K51.0** \_\_\_\_\_ Universal Ulcerative (chronic) Pancolitis
- K51.9** \_\_\_\_\_ Ulcerative Colitis, Unspecified
- K60.3** Anal Fistula
- K63.2** Fistula of Intestine
- Other **ICD-10 Code** (Diagnosis/Description): \_\_\_\_\_

**4. Pre-medications:**  None **OR** Administered 30 minutes prior to infusion as selected:

*\*Product information suggests premedication of antihistamines, acetaminophen, and/or corticosteroids.*

Acetaminophen: <input type="checkbox"/> 650mgs PO <input type="checkbox"/> 500mgs PO <input type="checkbox"/> 325mgs PO	Diphenhydramine: <input type="checkbox"/> 25 mgs PO, <input type="checkbox"/> 50mgs PO, <input type="checkbox"/> 25 mgs IVP, <input type="checkbox"/> 50mgs IVP or Alternate oral antihistamine to diphenhydramine: <input type="checkbox"/> Cetirizine 10 mg, <input type="checkbox"/> Loratadine 10 mg, Fexofenadine <input type="checkbox"/> 60mgs or <input type="checkbox"/> 180mgs Methylprednisolone <input type="checkbox"/> 40mgs IVP <input type="checkbox"/> 125mgs IVP or other _____mgs IVP Famotidine: <input type="checkbox"/> 20mgs PO, <input type="checkbox"/> 40mgs PO, <input type="checkbox"/> 20mgs IVP, <input type="checkbox"/> 40mgs IVP
--	---

Pre-medicate with other: \_\_\_\_\_

*CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion*

**5. Dose:** RENFLEXIS (infliximab-abda) infused via pump with 0.22 micron filter  
 \_\_\_\_\_ 5 mg/kg per 250 ml Sodium Chloride 0.9% IV to infuse over at least 2 hours **OR**  
 \_\_\_\_\_ Other dose: \_\_\_\_\_ mg or \_\_\_\_\_ mg/kg per 250 - 500 ml Sodium Chloride 0.9% IV

**6. Frequency:**  
 \_\_\_\_\_ Induction orders to be completed at 0 week, 2 week, and 6 weeks, and then every 8 weeks.  
 \_\_\_\_\_ Orders every 8 weeks (maintenance).

**Special orders:** \_\_\_\_\_

***If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES***

**7. Physician's Signature:** \_\_\_\_\_ / \_\_\_\_\_ Date: \_\_\_\_\_  
 No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: \_\_\_\_\_ NPI: \_\_\_\_\_

**8. Fax updated supporting clinical MD notes with each order renewal or change in orders**  
*Infusion order forms and Adverse Drug Reaction Guidelines are available at [www.palmettoinfusion.com](http://www.palmettoinfusion.com)*



MRN: \_\_\_\_\_

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: \_\_\_\_\_

## Guidelines for Prescribing RENFLEXIS (infliximab-abda) for Gastroenterology (Required documentation with all initial referrals)

Patient Name: \_\_\_\_\_

Referral Date: \_\_\_\_\_

\_\_\_\_\_ Include signed and completed **Plan of Treatment**. (*MD must complete sections 1-8*)  
(Infusion order forms & Standard Adverse Reactions orders are available at [www.palmettoinfusion.com](http://www.palmettoinfusion.com) under Agency/MD tab)

\_\_\_\_\_ Include patient demographic information and insurance information. (Copy of insurance cards if available)

\_\_\_\_\_ Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.

**INFLECTRA® is a biosimilar to Remicade® that is indications for:**

- **Adult Crohn's Disease:** reduces signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active disease who have had an **inadequate response to conventional therapy**. Reduces the number of draining enterocutaneous/rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing disease. **At a dose of 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Some adult patients who initially respond to treatment may benefit from increasing the dose to 10 mg/kg if they later lose their response.**
- **Pediatric Crohn's Disease:** reduces signs/symptoms and inducing/maintaining clinical remission in pediatric patients with moderately to severely active disease who have had an **inadequate response to conventional therapy**. **At a dose of 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks.**
- **Ulcerative Colitis:** reduces signs/symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in adult patients with moderately to severely active disease who have had an **inadequate response to conventional therapy**. **At a dose of 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks.**

\_\_\_\_\_ If patient is switching biological therapies, then MD must specify wash-out period prior to starting RENFLEXIS as specified of \_\_\_\_\_ weeks. Last known biological therapy: \_\_\_\_\_ and last date received: \_\_\_\_\_. (Include copy of last infusion record if available and currently on therapy)

\_\_\_\_\_ Other as requested: \_\_\_\_\_

**Pre-Screening: (TB and Hepatitis screening results must be available prior to start of therapy and within last 12 months.)**

\_\_\_\_\_ **Required TB screening results: PPD or QuantiFERON Gold Test.**  
(\* If screening results are positive or indeterminate, then a negative CXR result is required.)

\_\_\_\_\_ **Required Hepatitis screening to include: Hepatitis B Surface Antigen results.**

**\*\* Warnings/Precautions: \*Product information suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of infliximab or biosimilar.** Patient should not have an active ongoing infection, signs or symptoms of malignancy, or invasive fungal infection. Do not initiate RENFLEXIS therapy in patients with moderate to severe Congestive Heart Failure. **RENFLEXIS at doses of >5 mg/kg should not be administered to patients with moderate to severe heart failure.** Patient with mild CHF should be closely monitored. Therapy should be discontinued in patients who develop new or worsening symptoms of heart failure. **Hepatotoxicity:** Stop therapy in case of jaundice and/or marked liver enzyme elevations. **Malignancies:** including lymphoma was greater in TNF blocker treated patients Due to the risk of HSTCL carefully assess the risk/benefits especially if the patient has Crohn's disease or ulcerative colitis, is male, and is receiving azathioprine or 6-mercaptopurine treatment. **Lupus-like syndrome:** stop if symptoms develop. Evaluation of immunizations should be completed prior to and live vaccines should not be given before or concurrently with RENFLEXIS. See full prescribing information.

**Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.**

**Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.**