Palmetto INFUSION

Phone: 800-809-1265

MRN:_____
DOB:_____

Standard Tysabri Plan of Treatment

NOTE: Patient ***may be ineligible*** to receive Tysabri if receiving antibiotic for active infectious process (due to the possibility of developing a super-infection related to its effect on the immune status), or if he/she has a suspected infection.

Name: _____Height_____ Weight:

Allergies:_____

Primary Diagnosis: **ICD-10 G35 Relapsing Multiple Sclerosis**

This section must be completed by the referring physician.

Patient has been on the following medication(s) for treatment of MS (Must have failed 2 or more MS medications):
_____.

These medications will be discontinued as of_____.

Patient may receive Tysabri after wash-out period of_____ weeks or_____months.

If adverse drug reaction, Implement the Standing Adverse Reaction protocol.

Premedicate 30 minutes prior to infusion with 650 mg Acetaminophen PO and one of the following *oral* antihistamines:

☐ Diphenhydramine 50mg ☐ Fexofenadine 60mg ☐ Fexofenadine 180mg ☐ Cetirizine10mg ☐ Loratadine 10 mg

OR Premedicate with other _____

Drug:

Tysabri 300mg in 100ml NS IV over 1 hour Every 4 weeks, Monitor pt. for 1 hour post infusion

Pharmacist to perform clinical drug monitoring.

(No Stamped Signatures please)

Physician's Signature:_____/_____Date: _____

(Dispense as written) (Substitution permitted)

Print Physician Name:_____

PLEASE FAX DEMOGRAPHICS AND INSURANCE INFORMATION TO:

866-872-8920

Revised 8/28/20