



MRN: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: _____

STANDARD INFLECTRA® (infliximab-dyyb) PLAN OF TREATMENT FOR GASTROENTEROLOGY

NOTE: Patient *may be ineligible* to receive (infliximab-dyyb) if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

1. Patient Name: _____ Height (inches): _____ Weight (lbs): _____

2. Allergies: _____

3. DIAGNOSIS: * Please complete the 2nd and 3rd digits to complete the ICD-10 code for billing

- K50.0** _____ Crohn’s Disease (small intestine) **K51.8** _____ Other Ulcerative (chronic) Colitis
- K50.1** _____ Crohn’s Disease (large intestine) **K51.5** _____ Left sided Ulcerative (chronic) Colitis
- K50.8** _____ Crohn’s Disease (small & large intestine) **K51.0** _____ Universal Ulcerative (chronic) Pancolitis
- K50.9** _____ Crohn’s Disease, Unspecified **K51.9** _____ Ulcerative Colitis, Unspecified **K60.3** Anal Fistula
- K63.2** Fistula of Intestine Other **ICD-10 Code** (Diagnosis/Description): _____

4. Pre-medications: Administered 30 minutes prior to infusion as selected:

**Product information suggests premedication of antihistamines, acetaminophen, and/or corticosteroids.*

<p>a) Acetaminophen:</p> <ul style="list-style-type: none"> <input type="checkbox"/> 650mgs PO <input type="checkbox"/> 500mgs PO <input type="checkbox"/> 325mgs PO 	<p>b) Diphenhydramine: <input type="checkbox"/> 25 mgs PO, <input type="checkbox"/> 50mgs PO, <input type="checkbox"/> 25 mgs IVP, <input type="checkbox"/> 50mgs IVP or</p> <p>c) Alternate oral antihistamine to diphenhydramine:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Cetirizine 10 mg, <input type="checkbox"/> Loratadine 10 mg, Fexofenadine <input type="checkbox"/> 60mgs or <input type="checkbox"/> 180mgs <p>d) Other: Methylprednisolone <input type="checkbox"/> 40mgs IVP <input type="checkbox"/> 125mgs IVP or other _____mgs IVP Famotidine: <input type="checkbox"/> 20mgs PO, <input type="checkbox"/> 40mgs PO, <input type="checkbox"/> 20mgs IVP, <input type="checkbox"/> 40mgs IVP</p>
---	---

e) Pre-medicate with other: _____

CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion

If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES.

5. Order:

INFLECTRA® (infliximab-dyyb) 5 mg/kg per 250 ml Sodium Chloride 0.9% IV to infuse over at least 2 hours

OR

Other Dose: _____ mg or _____ mg/kg per 250 - 500 ml Sodium Chloride 0.9% IV

- Induction orders to be completed at 0 week, 2 week, and 6 weeks
- Maintenance Orders every 8 weeks

Special Orders: _____

Lab orders with infusions: _____

7. Physician’s Signature: _____ / _____ Date: _____
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician’s Name with Credentials: _____

8. Fax updated supporting clinical MD notes with each order renewal or change in orders

Infusion order forms available at www.palmettoinfusion.com



MRN: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: _____

Guidelines for Prescribing INFLECTRA® (infliximab-dyyb) for Gastroenterology

(Required documentation with all initial referrals)

Patient Name: _____

Referral Date: _____

_____ Include signed and completed **Plan of Treatment**. (*MD must complete sections 1-8*)
 (*Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab*)

_____ Include patient demographic information and insurance information. (Copy of insurance cards if available)

_____ **Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.**

INFLECTRA® is a biosimilar to Remicade® that is indications for:

- **Adult Crohn's Disease:** reduces signs and symptoms and inducing and maintaining clinical remission in adult patients with moderately to severely active disease who have had an **inadequate response to conventional therapy**. Reduces the number of draining enterocutaneous/rectovaginal fistulas and maintaining fistula closure in adult patients with fistulizing disease. **At a dose of 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Some adult patients who initially respond to treatment may benefit from increasing the dose to 10 mg/kg if they later lose their response.**
- **Pediatric Crohn's Disease:** reduces signs/symptoms and inducing/maintaining clinical remission in pediatric patients with moderately to severely active disease who have had an **inadequate response to conventional therapy**. **At a dose of 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks. Pediatric Use –INFLECTRA® has not been studied in children with Crohn's disease or ulcerative colitis.**
- **Ulcerative Colitis:** reduces signs/symptoms, inducing and maintaining clinical remission and mucosal healing, and eliminating corticosteroid use in adult patients with moderately to severely active disease who have had an **inadequate response to conventional therapy**. **At a dose of 5 mg/kg at 0, 2 and 6 weeks, then every 8 weeks.**

_____ If patient is switching biological therapies, then MD must specify wash-out period prior to starting INFLECTRA® as specified of _____ weeks. Last known biological therapy: _____ and last date received: _____. (Include copy of last INFLECTRA® infusion record if available and currently on therapy)

_____ Other as requested: _____

Pre-Screening: (TB and Hepatitis screening results must be available prior to start of therapy and within last 12 months.)

_____ **Required TB screening results: PPD or QuantiFERON Gold Test.**
 (* If screening results are positive or indeterminate, then a negative CXR result is required.)

_____ **Required Hepatitis screening to include: Hepatitis B Surface Antigen results.**

**** Warnings/Precautions: *Product information suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of infliximab or biosimilar.** Patient should not have an active ongoing infection, signs or symptoms of malignancy, or invasive fungal infection. Do not initiate INFLECTRA® therapy in patients with moderate to severe Congestive Heart Failure. **INFLECTRA® at doses of >5 mg/kg should not be administer to patients with moderate to severe heart failure.** Patient with mild CHF should be closely monitored. Therapy should be discontinued in patients who develop new or worsening symptoms of heart failure. **Hepatotoxicity:** Stop therapy in case of jaundice and/or marked liver enzyme elevations. Evaluation of immunizations should be completed prior to and live vaccines should not be given before or concurrently with INFLECTRA®. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.