





MRN: \_\_\_\_\_

Phone: 1-800-809-1265 Fax: 1-866-872-8920

DOB: \_\_\_\_\_

### Guidelines for Prescribing Truxima® (rituximab-abbs) for GPA or MPA (Required documentation with all initial referrals)

Patient Name: \_\_\_\_\_

Referral Date: \_\_\_\_\_

\_\_\_ Include signed and completed **Plan of Treatment**. (MD must complete sections 1-8)  
(Infusion order forms & Standard Adverse Reactions orders are available at [www.palmettoinfusion.com](http://www.palmettoinfusion.com) under Agency/MD tab)

\_\_\_ Include patient demographic information and insurance information. (Copy of insurance cards if available)

\_\_\_ **Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.**

\* Rituximab-abbs, in combination with glucocorticoids, is indicated for the treatment of adult patients with Granulomatosis with Polyangiitis (GPA) (Wegener’s Granulomatosis) and Microscopic Polyangiitis (MPA).

\_\_\_ If patient is switching biological therapies, then MD must specify wash-out period prior to starting Truxima® as specified of \_\_\_\_\_ weeks. Last known biological therapy: \_\_\_\_\_ and last date received: \_\_\_\_\_.

\_\_\_ Other as requested: \_\_\_\_\_

#### Pre-Screening: (Hepatitis screening results must be available prior to start of therapy and within last 12 months)

\_\_\_ **Required Hepatitis screening to include: Hepatitis B Surface Antigen (HBsAg) and Total Hepatitis B Core Antibody (anti- Hbc)**

\*Truxima® is contraindicated in patient with active HBV. Patients who are negative for surface antigen HBsAg (-) and positive for HB core antibody HbcAB (+) or positive for surface antigen HBsAg (+), should consult liver disease experts before starting and during treatment.

\***Other:** Glucocorticoids administered as methylprednisolone 1000 mg intravenously daily for 1 to 3 days followed by oral prednisone 1 mg/kg/day (not to exceed 80 mg/day and tapered per clinical need) are recommended to treat severe vasculitis symptoms. This regimen should begin within 14 days prior to or with the initiation of rituximab and may continue during and after the 4-week course of rituximab treatment. Methylprednisolone Plan of Treatment are available at [www.palmettoinfusion.com](http://www.palmettoinfusion.com) if needed.

\*\* Warnings/Precautions: • **Hepatitis B Virus Reactivation**- Screen all patients for HBV infection by measuring HBsAg and anti-HBc (antibodies) before initiating treatment with Truxima®. For patients who show evidence of prior hepatitis B infection (HBsAg positive or HBsAg negative but anti-HBc positive), consult with physicians with expertise in managing hepatitis B regarding monitoring and consideration for HBV antiviral therapy before and/or during Truxima® treatment. HBV reactivation has been reported up to 24 months following completion of Rituxan® •Glucocorticoids administered methylprednisolone IV premed or its equivalent 30 minutes prior to each infusion are recommended to reduce the incidence and severity of infusion reactions. Evaluation of immunizations should be completed prior to and live vaccines should not be given before or concurrently. **Serious Infections:** including fatal, bacterial, fungal, and new or reactivated viral infections can occur during and following the completion of Truxima®. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient’s insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.