



MRN: _____
 DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

STANDARD INFLECTRA® (infliximab-dyyb) PLAN OF TREATMENT FOR DERMATOLOGY

NOTE: Patient *may be ineligible* to receive (infliximab-dyyb) if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

1. **Patient Name:** _____ Height (inches): _____ Weight (lbs): _____

2. **Allergies:** _____

3. **DIAGNOSIS:** * Please complete the 2nd and 3rd digits to complete the ICD-10 code for billing

L40.5 _____ Psoriatic Arthritis/Arthropathy L40. _____ Psoriasis

Other **ICD-10 Code:** _____ **Diagnosis description:** _____

4. **Pre-medications: Administered 30 minutes prior to infusion as selected:**

**Product information suggests premedication of antihistamines, acetaminophen, and/or corticosteroids.*

| | |
|---|---|
| a) Acetaminophen: <input type="checkbox"/> 650mgs PO <input type="checkbox"/> 500mgs PO <input type="checkbox"/> 325mgs PO | b) Diphenhydramine: <input type="checkbox"/> 25 mgs PO, <input type="checkbox"/> 50mgs PO, <input type="checkbox"/> 25 mgs IVP, <input type="checkbox"/> 50mgs IVP or c) Alternate oral antihistamine to diphenhydramine: <input type="checkbox"/> Cetirizine 10 mg, <input type="checkbox"/> Loratadine 10 mg, Fexofenadine <input type="checkbox"/> 60mgs or <input type="checkbox"/> 180mgs d) Other: Methylprednisolone <input type="checkbox"/> 40mgs IVP <input type="checkbox"/> 125mgs IVP or other _____mgs IVP Famotidine: <input type="checkbox"/> 20mgs PO, <input type="checkbox"/> 40mgs PO, <input type="checkbox"/> 20mgs IVP, <input type="checkbox"/> 40mgs IVP |
|---|---|

e) Pre-medicate with other: _____

CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion

Orders: Obtain weight each visit. **Vital signs at baseline, every 30 minutes beginning with start of infusion, at completion, 30 minutes after completion for the first 3 treatments, and then may discharge when infusion is complete.** Instruct patient/caregiver on medications and signs/symptoms of adverse reaction. Assess patient for response to therapy. Utilize existing central line for administration or initiate a peripheral IV with each infusion as needed. Sodium Chloride 0.9% flush 3-10 ml before, after, and as needed during the infusion. Follow infusion with Heparin 100 units/ml 1 – 5 ml per line type or to peripheral IV as required for multiple day treatments. Pump, tubing, 0.22 micron filter, and supplies needed to complete prescribed therapy. Pharmacist to perform clinical drug monitoring. ***If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES.***

5. **Dose:** _____ INFLECTRA® (infliximab-dyyb) 5 mg/kg per 250 ml Sodium Chloride 0.9% IV to infuse over at least 2 hours **OR**
 _____ Other Dose: _____ mg or _____ mg/kg per 250 - 500 ml Sodium Chloride 0.9% IV

6. **Frequency:** _____ Induction orders to be completed at 0 week, 2 week, and 6 week
 _____ Maintenance Orders every 8 weeks
 _____ Special Orders: _____

Lab orders with infusions: _____

7. **Physician's Signature:** _____ / _____ Date: _____
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: _____

8. Fax updated supporting clinical MD notes with each order renewal or change in orders
Infusion order forms available at www.palmettoinfusion.com



MRN: _____

DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Guidelines for Prescribing INFLECTRA® (infliximab-dyyb) for Dermatology (Required documentation with all initial referrals)

Patient Name: _____

Referral Date: _____

___ Include signed and completed **Plan of Treatment**. (MD must complete sections 1-8)
(Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab)

___ Include patient demographic information and insurance information. (Copy of insurance cards if

___ **Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.**

INFLECTRA® is a biosimilar to Remicade® that is indications for:

- **Psoriatic Arthritis & Plaque Psoriasis:** indicated for dosing at 5 mg/kg induction & then every 8 weeks.

___ If patient is switching biological therapies, then MD must specify wash-out period prior to starting INFLECTRA® as specified of _____ weeks. Last known biological therapy: _____ and last date received: _____. (Include copy of last INFLECTRA® infusion record if available and currently on therapy)

___ Other as requested: _____

Pre-Screening:

___ **Required TB screening results: PPD or QuantiFERON Gold Test.**
(* If screening results are positive or indeterminate, then a negative CXR result is required.)

___ **Required Hepatitis screening to include: Hepatitis B Surface Antigen results.**

**** Warnings/Precautions: *Product information suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of infliximab or biosimilar.** Patient should not have an active ongoing infection, signs or symptoms of malignancy, or invasive fungal infection. Do not initiate INFLECTRA® therapy in patients with moderate to severe Congestive Heart Failure. **INFLECTRA® at doses of >5 mg/kg should not be administer to patients with moderate to severe heart failure.** Patient with mild CHF should be closely monitored. Therapy should be discontinued in patients who develop new or worsening symptoms of heart failure. **Hepatotoxicity:** Stop therapy in case of jaundice and/or marked liver enzyme elevations. Evaluation of immunizations should be completed prior to and live vaccines should not be given before or concurrently with INFLECTRA®. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.