

CONFIDENTIAL Property of Palmetto Infusion Services / CONFIDENTIAL Property of Palmetto Infusion Services / CONFIDENTIAL Property of Palmetto Infusion Services

Phone: 1-800-809-1265 Fax: 1-866-872-8920

7. Physician's Signature: ______
No Stamp Signatures

MRN:		
· -		
DOB.		

STANDARD RENFLEXIS (infliximab-abda) PLAN OF TREATMENT NOTE: Patient may be ineligible to receive (infliximab-abda) if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery. 1. Patient Name: ______ Height (inches): _____ Weight (lbs): _____ 2. Allergies: 3. Diagnosis: ICD-10 Code: Diagnosis Description: 4. Pre-medications: Administered 30 minutes prior to infusion as selected: *Product information suggests premedication of antihistamines, acetaminophen, and/or corticosteroids. Diphenhydramine: □ 25 mgs PO, □ 50mgs PO, □ 25 mgs IVP, □ 50mgs IVP or Acetaminophen: □ 650mgs PO Alternate oral antihistamine to diphenhydramine: □ 500mgs PO □ Cetirizine 10 mg, □ Loratadine 10 mg, Fexofenadine □ 60mgs or □ 180mgs □ 325mgs PO Methylprednisolone

□ 40mgs IVP □ 125mgs IVP or other mgs IVP Famotidine: □ 20mgs PO, □ 40mgs PO, □ 20mgs IVP, □ 40mgs IVP Pre-medicate with other: CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion **5. Dose**: RENFLEXIS (infliximab-abda) infused via pump with 0.22 micron filter 3 mg/kg per 250 ml Sodium Chloride 0.9% IV to infuse over at least 2 hours OR Other Dose: _____mg or ____mg/kg per 250 - 500 ml Sodium Chloride 0.9% IV 6. Frequency: _____ Induction orders to be completed at 0 week, 2 week, and 6 weeks, and then every 8 weeks Orders every 8 weeks (maintenance) Special orders: ______ If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES

8. Fax updated supporting clinical MD notes with each order renewal or change in ordersInfusion order forms and Adverse Drug Reaction Guidelines are available at www.palmettoinfusion.com

(Substitution permitted)

(Dispense as written)

Printed Physician's Name with Credentials:______



Phone: 1-800-809-1265

MKN:_			
DOB.			

Standard Guidelines for Prescribing RENFLEXIS (infliximab-abda)

(Required documentation with all initial referrals)

Patien	t Name: Referral Date:
	Include signed and completed Plan of Treatment . (MD must complete sections 1-8) (Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab)
	Include patient demographic information and insurance information. (Copy of insurance cards if available)
	Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.
	If patient is switching biological therapies, then MD must specify wash-out period prior to starting RENFLEXIS as specified of weeks. Last known biological therapy: and last date received: (Include copy of last infusion record if available and currently on therapy)
	Other as requested:
Requi	red Pre-Screening:
	TB screening test completed. Results (positive or negative):
	Hepatitis B screening test completed. Results (positive or negative):
	*If TB or Hep B results are positive, please send documentation of treatment or medical clearance.

** Warnings/Precautions: *Product information suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of infliximab or biosimilar. Patient should not have an active ongoing infection, signs or symptoms of malignancy, or invasive fungal infection. Do not initiate RENFLEXIS therapy in patients with moderate to severe Congestive Heart Failure. RENFLEXIS at doses of >5 mg/kg should not be administered to patients with moderate to severe heart failure. Patient with mild CHF should be closely monitored. Therapy should be discontinued in patients who develop new or worsening symptoms of heart failure. Hepatotoxicity: Stop therapy in case of jaundice and/or marked liver enzyme elevations. Malignancies: including lymphoma was greater in TNF blocker treated patients Due to the risk of HSTCL carefully assess the risk/benefits especially if the patient has Crohn's disease or ulcerative colitis, is male, and is receiving azathioprine or 6-mercaptopurine treatment. Lupus-like syndrome: stop if symptoms develop. Evaluation of immunizations should be completed prior to and live vaccines should not be given before or concurrently with RENFLEXIS. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.