



MRN: _____

DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

STANDARD Xolair® (omalizumab) PLAN OF TREATMENT FOR NASAL POLYPS

NOTE: Patient *may be ineligible* to receive Xolair® (omalizumab) if patient has signs/symptoms of parasitic infection, is currently being treated for a parasitic infection, or is having acute bronchospasm and/or asthma attack.

1. Patient Name: _____ Height (inches): _____ Weight (lbs): _____

2. Allergies: _____

3. Diagnosis: J33.0 Polyp of nasal cavity J33.1 Polypoid sinus degeneration
 J33.8 Other polyp of sinus J33.9 Nasal polyp, unspecified
 Other ICD-10 Code/description: _____

CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion / CONFIDENTIAL Property of Palmetto Infusion

4. Dose/Frequency:

Xolair® (omalizumab) subcutaneously every 2 weeks: 300 mg/dose 375 mg/dose 450 mg/dose
 525 mg/dose 600mg/dose
Xolair® (omalizumab) subcutaneously every 4 weeks: 75 mg/dose 150 mg/dose 225mg/dose
 300mg/dose 450mg/dose 600 mg/dose

Special Orders: _____
(Administer as subcutaneous injection to upper arm, thigh, or abdomen. No more than 150 mg per injection site)

Extended post treatment monitoring for any patient new to therapy: monitor patient for two (2) hours after first injection, for one (1) hour after second injection, for 30-minutes after third injection, and then all patients for 15-minutes after each injection thereafter.

If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES

5. Physician's Signature: _____ / _____ Date: _____
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: _____ NPI: _____

6. Fax updated supporting clinical MD notes with each order renewal or change in orders
Infusion order forms and Adverse Drug Reaction Guidelines are available at www.palmettoinfusion.com



MRN: _____

DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Guidelines for Prescribing Xolair® (omalizumab)

(Required documentation with all initial referrals)

Patient Name: _____

Referral Date: _____

_____ Include signed and completed **Plan of Treatment**. (MD must complete sections 1-6)
(Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab)

_____ Include patient demographic information and insurance information. (Copy of insurance cards if available)

Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any documented use of inhaled or oral corticosteroids, leukotriene modifiers, short or long acting beta-agonists. Any lab results, Pulmonary Function Tests (FEV₁ and Peak Flow), and/or positive skin test or RAST testing to support diagnosis.

Xolair® is an anti-IgE antibody indicated for add-on maintenance treatment of **nasal polyps** in adult patients 18 years of age and older with inadequate response to nasal corticosteroids. Administer XOLAIR 75mg to 600mg by subcutaneous injection every 2 or 4 weeks based on serum total IgE level (IU/mL) measure before the start of treatment and by body weight (kg) [see Prescribing Information Dosage and Administration 2.2].

If patient is switching biologic therapies, then MD must specify wash-out period prior to starting Xolair® as specified of _____ weeks. Last known therapy: _____ & last known date received: _____.

Other as requested: _____

Pre-Screening:

Pre-treatment serum IgE level as required for dosing for nasal polyps. Dosage is determined by initial IgE level and body weight. Do NOT use levels for subsequent dose determinations unless treatment has been interrupted for more than 1 year.

**** Warnings/Precautions: Anaphylaxis reactions:** such as bronchospasm, hypotension, syncope, urticaria, and/or angioedema of the throat or tongue, has been reported to occur after administration of Xolair®. Anaphylaxis has occurred as early as after the first dose of Xolair®, but also has occurred beyond 1 year after beginning regularly administered treatment. Fever, Arthralgia, and Rash: Stop Xolair® if patients develop signs and symptoms similar to serum sickness. **Acute Asthma Symptoms:** Do not use for the treatment of acute bronchospasm or status asthmaticus. **Malignancy:** have been observed in clinical studies. **Corticosteroid Reduction:** Do not abruptly discontinue corticosteroids upon initiation of therapy. **Eosinophilic Conditions:** Be alert to eosinophilia, vasculitic rash, worsening pulmonary symptoms, cardiac complications, and/or neuropathy, especially upon reduction of oral corticosteroids. **Helminth Infections:** Treat patients with pre-existing parasitic infections before therapy. If patients become infected while receiving treatment and do not respond to anti-helminth treatment, then discontinue Xolair® until parasitic infection resolves. **Pregnancy/Breastfeeding:** Discuss Pregnancy or breastfeeding plans/risks prior to start of therapy. See full prescribing information.

Palmetto Infusion Services will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.