

MRN: \_\_\_\_\_

DOB: \_\_\_\_\_

### STANDARD REMICADE® (infliximab) PLAN OF TREATMENT FOR GASTROENTEROLOGY

**NOTE:** Patient *may be ineligible* to receive infliximab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

**1. Patient Name:** \_\_\_\_\_ Height (inches): \_\_\_\_\_ Weight (lbs): \_\_\_\_\_

**2. Allergies:** \_\_\_\_\_

**3. DIAGNOSIS:** \* Please complete the 2<sup>nd</sup> and 3<sup>rd</sup> digits to complete the ICD-10 code for billing

- |   |   |  |
|---|---|--|
| <input type="checkbox"/> <b>K50.0</b> _____ Crohn's Disease (small intestine)   | <input type="checkbox"/> <b>K51.8</b> _____ Other Ulcerative (chronic) Colitis        |  |
| <input type="checkbox"/> <b>K50.1</b> _____ Crohn's Disease (large intestine)   | <input type="checkbox"/> <b>K51.5</b> _____ Left sided Ulcerative (chronic) Colitis   |  |
| <input type="checkbox"/> <b>K50.8</b> _____ Crohn's Disease (small & large intestine)   | <input type="checkbox"/> <b>K51.0</b> _____ Universal Ulcerative (chronic) Pancolitis |  |
| <input type="checkbox"/> <b>K50.9</b> _____ Crohn's Disease, Unspecified  | <input type="checkbox"/> <b>K51.9</b> _____ Ulcerative Colitis, Unspecified           | <input type="checkbox"/> <b>K60.3</b> Anal Fistula |
| <input type="checkbox"/> <b>K63.2</b> Fistula of Intestine <input type="checkbox"/> Other <b>ICD-10 Code</b> (Diagnosis/Description): _____ |   |  |

**4. Pre-medications: Administered 30 minutes prior to infusion as selected:**

*\*Product information suggests premedication of antihistamines, acetaminophen, and/or corticosteroids.*

a) Acetaminophen: <input type="checkbox"/> 650mgs PO <input type="checkbox"/> 500mgs PO <input type="checkbox"/> 325mgs PO	b) Diphenhydramine: <input type="checkbox"/> 25 mgs PO, <input type="checkbox"/> 50mgs PO, <input type="checkbox"/> 25 mgs IVP, <input type="checkbox"/> 50mgs IVP or c) Alternate oral antihistamine to diphenhydramine: <input type="checkbox"/> Cetirizine 10 mg, <input type="checkbox"/> Loratadine 10 mg, Fexofenadine <input type="checkbox"/> 60mgs or <input type="checkbox"/> 180mgs d) Other: Methylprednisolone <input type="checkbox"/> 40mgs IVP <input type="checkbox"/> 125mgs IVP or other _____ mgs IVP Famotidine: <input type="checkbox"/> 20mgs PO, <input type="checkbox"/> 40mgs PO, <input type="checkbox"/> 20mgs IVP, <input type="checkbox"/> 40mgs IVP
---	--

e) Pre-medicate with other: \_\_\_\_\_

**5. Dose:**  Remicade® (infliximab) 5 mg/kg per 250 ml Sodium Chloride 0.9% IV to infuse over at least 2 hours **OR**  
 Other Dose: \_\_\_\_\_ mg or \_\_\_\_\_ mg/kg per 250 - 500 ml Sodium Chloride 0.9% IV

**6. Frequency:**  Induction orders to be completed at 0 week, 2 week, and 6 weeks, and then every 8 weeks thereafter  
 Orders every 8 weeks (maintenance).  
 Special Orders: \_\_\_\_\_

Lab orders with infusions: \_\_\_\_\_

**7. Physician's Signature:** \_\_\_\_\_ / \_\_\_\_\_ Date: \_\_\_\_\_  
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: \_\_\_\_\_

MRN: \_\_\_\_\_

DOB: \_\_\_\_\_

## Guidelines for Prescribing Remicade<sup>®</sup> (infliximab) for Gastroenterology (Required documentation with all initial referrals)

Patient Name: \_\_\_\_\_

Referral Date: \_\_\_\_\_

- Include signed and completed **Plan of Treatment**.
- Include patient demographic information and insurance information. (Copy of insurance cards if available)
- Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.**
  - Remicade<sup>®</sup> is indicated for **Adult Crohn's Disease**: Dose of 5 mg/kg induction & then every 8 weeks. For patients who respond and then lose response, consider treatment with 10mg/kg. Patients who do not respond by 14 weeks are unlikely to with continued dosing.
  - **Pediatric Crohn's Disease, Adult and Pediatric Ulcerative Colitis**: Dose of 5mg/kg induction and then every 8 weeks.
- If patient is switching biological therapies, then MD must specify wash-out period prior to starting Remicade<sup>®</sup> as specified of \_\_\_\_\_ weeks. Last known biological therapy: \_\_\_\_\_ and last date received: \_\_\_\_\_. (Include copy of last Remicade<sup>®</sup> infusion record if available and currently on therapy)
- Other as requested: \_\_\_\_\_

### Pre-Screening:

- Required TB screening results: PPD or QuantiFERON Gold Test.**  
(\* If screening results are positive or indeterminate, then a negative CXR result is required.)
- Required Hepatitis screening to include: Hepatitis B Surface Antigen results.**

*\*Product information suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of therapy.*

\*\* Warnings/Precautions: **Serious Infections**: Patient should not have an active ongoing infection, signs or symptoms of malignancy, or invasive fungal infection. Do not initiate Remicade<sup>®</sup> (infliximab) therapy in patients with **moderate to severe Congestive Heart Failure**. **Remicade<sup>®</sup> (infliximab) at doses of >5 mg/kg should not be administer to patients with moderate to severe heart failure.** Patient with mild CHF currently receiving Remicade<sup>®</sup> (infliximab) should be closely monitored. Therapy should be discontinued in patients who develop new or worsening symptoms of heart failure. Evaluation of immunizations should be completed prior to and live vaccines should not be given before or concurrently with Remicade<sup>®</sup>. See full prescribing information.



Checklist for referrals to AccuRX Infusion Center:

Fax referral to 1-866-990-3192

- Patient demographics – address, phone number, SS#, etc.
- Insurance Information – copy of the card(s) if possible
- Plan of Treatment/Orders
- Most recent physician office notes to include tried and failed therapies – all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs
- Any lab results or other diagnostic procedures to support the diagnosis

AccuRx Infusion Center will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.