

MRN: \_\_\_\_\_

DOB: \_\_\_\_\_

## STANDARD Tysabri® (natalizumab) PLAN OF TREATMENT for MULTIPLE SCLEROSIS

**NOTE:** Patient **may be ineligible** to receive natalizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new-onset or deterioration neurological changes, and/or surgery.

**1. Patient Name:** \_\_\_\_\_ Height (inches): \_\_\_\_\_ Weight (lbs): \_\_\_\_\_

**2. Allergies:** \_\_\_\_\_

**3. Primary Diagnosis: G35 Relapsing Multiple Sclerosis**

**4. Pre-medications: Administered 30 minutes prior to infusion as selected:**

*\*Product information suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of therapy.*

Acetaminophen: <input type="checkbox"/> 650 mg PO <input type="checkbox"/> 500 mg PO <input type="checkbox"/> 325 mg PO	Diphenhydramine: <input type="checkbox"/> 25 mg PO, <input type="checkbox"/> 50 mg PO, <input type="checkbox"/> 25 mg IVP, <input type="checkbox"/> 50 mg IVP or Fexofenadine <input type="checkbox"/> 60 mg or <input type="checkbox"/> 180 mg, <input type="checkbox"/> Cetirizine 10 mg, <input type="checkbox"/> Loratadine 10 mg Methylprednisolone <input type="checkbox"/> 40 mg IVP <input type="checkbox"/> 125 mg IVP or other _____ mg IVP Famotidine: <input type="checkbox"/> 20 mg PO, <input type="checkbox"/> 40 mg PO, <input type="checkbox"/> 20 mg IVP, <input type="checkbox"/> 40 mg IVP
--	---

Pre-medicate with other: \_\_\_\_\_

### Dose:

Tysabri® (natalizumab) 300 mg per 100 ml Sodium Chloride 0.9% IV to infuse over at least 1- hour with no filter required. Follow each infusion with 1-hour post infusion monitoring.

**Frequency:** Dosing every 4 weeks, no less than every 28 days.

**\*Prior to each infusion:** ensure that the patient has a current Notice of Patient Authorization on file to receive Tysabri® (natalizumab) for Multiple Sclerosis and complete/submit Pre-infusion Patient Checklist within 24 hours to Biogen Idec.

**Special orders:** \_\_\_\_\_

***If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES***

**5. Physician's Signature:** \_\_\_\_\_ / \_\_\_\_\_ Date: \_\_\_\_\_  
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: \_\_\_\_\_ NPI: \_\_\_\_\_

MRN: \_\_\_\_\_

DOB: \_\_\_\_\_

## Guidelines for Prescribing Tysabri® (natalizumab) for Multiple Sclerosis (Required documentation with all initial referrals)

Patient Name: \_\_\_\_\_

Referral Date: \_\_\_\_\_

\_\_\_ Tysabri® (natalizumab) is restricted to credentialed prescribers and patients enrolled in the TOUCH® Prescribing Program. Contact TOUCH® Prescribing Program at 1-800-456-2255 for details and enrollment. TOUCH® Authorization form must be received prior to scheduling.

\_\_\_ Include signed and completed **Plan of Treatment**.

\_\_\_ Include patient demographic information and insurance information. (Copy of insurance cards if available)

\_\_\_ **Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any lab results and/or tests to support diagnosis.**

- TYSABRI® (natalizumab) is an integrin receptor antagonist indicated as monotherapy for the treatment of patients with relapsing forms of multiple sclerosis. TYSABRI increases the risk of progressive multifocal leukoencephalopathy (PML).

\_\_\_ Include the following:

Past tried and/or failed therapies: \_\_\_\_\_

Last known therapy: \_\_\_\_\_ and last date received: \_\_\_\_\_.

MD must specify wash-out period prior to starting Tysabri® as specified of \_\_\_\_\_ weeks.

\_\_\_ Other as requested: \_\_\_\_\_

### Pre-Screening:

\_\_\_ **anti-JCV antibodies test results within last 6 months.** (Patients who are anti-JCV antibody positive, will require documentation from referring MD that risks and benefits have been discussed)

\*\* Warnings/Precautions: TYSABRI® increases the risk of **progressive multifocal leukoencephalopathy (PML)**: an opportunistic viral infection of the brain that usually leads to death or severe disability • Risk factors for the development of PML include duration of therapy, prior use of immunosuppressants, and presence of anti-JCV antibodies. • Healthcare professionals should monitor patients on TYSABRI® for any new sign or symptom that may be suggestive of PML. TYSABRI® dosing should be withheld immediately at the first sign or symptom suggestive of PML. • **Herpes encephalitis and meningitis**: Life-threatening and fatal cases have occurred, discontinue TYSABRI® if this occurs • **Hepatotoxicity**: Significant liver injury, including liver failure requiring transplant, has occurred. Discontinue TYSABRI® in patients with evidence of liver injury. • **Immunosuppression/Infections**: TYSABRI® may increase the risk for certain infections including: pneumonias, urinary tract infections (including serious cases), gastroenteritis, vaginal infections, tooth infections, tonsillitis, and herpes infections. • **Hypersensitivity reactions**: Serious hypersensitivity reactions (e.g., anaphylaxis) have occurred. **Permanently discontinue TYSABRI® if such a reaction occurs.** • **Pregnancy Category C**. See full prescribing information.



Checklist for referrals to AccuRX Infusion Center:

Fax referral to 1-866-990-3192

- Patient demographics – address, phone number, SS#, etc.
- Insurance Information – copy of the card(s) if possible
- Plan of Treatment/Orders
- Most recent physician office notes to include tried and failed therapies – all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs
- Any lab results or other diagnostic procedures to support the diagnosis

AccuRx Infusion Center will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.