



3105 Sunset Boulevard
 West Columbia, SC 29169
 Phone: 803-771-7740
 Fax: 888-417-3658
 NPI 1265462659

Patient Name: _____ Diagnosis: _____

DOB: _____ MRN: _____

Rx

Medication: _____ Start Date: _____

Dose: _____ Concentration: _____ Route: _____

Frequency: _____ Duration: _____ Refills: _____

Labs: _____ Dispense: _____

- Palmetto Infusion RN to insert peripheral IV, monitor vital signs before, during and after infusion. May leave peripheral IV in place for 3 to 5 days if remains patent.
- Sodium Chloride 0.9% 5-10ml flush IV catheter/port before and after medication administration and/or as directed. Flush unused IV line(s) daily.
- Heparin 100units/ml **OR** Heparin 10 units/ml (usually for <18years)
1-5ml flush IV after Sodium Chloride 0.9% and/or as directed. Flush unused line(s) daily.
- Palmetto Infusion Services Pharmacist to manage above medication.**
- Pump, tubing, and supplies needed to complete prescribed therapy.
- Administer by CADD or Curlin Pump (Ambulatory Infusion Pump, Single or Multiple channels, Electric or Battery operated, with administrative equipment, worn by patient – E0781).
 - Dispense infusion supplies for external drug infusion pump, per cassette or bag (A4222).
 - Dispense supplies for maintenance of drug infusion catheter, per week (A4221).
- Administer by Freedom 60 Syringe Pump (Ambulatory Infusion Pump, Mechanical, Reusable, for subcutaneous infusion - E0779).
 - Dispense supplies for external drug infusion pump, syringe type cartridge, sterile, each (K0552).
- Palmetto Infusion RN to change vascular access device dressing weekly.
- Other: _____

_____ / _____ Date: _____

Dispense As Written

Substitution Permitted

Printed Physician name: _____ NPI: _____

Verbal order Read Back: _____ RPh/CPhT