

Phone: 1-800-809-1265 Fax: 1-866-872-8920

MRN: _____

DOB: _____

STANDARD Nucala® (mepolizumab) PLAN OF TREATMENT FOR ASTHMA

NOTE: Patient *may be ineligible* to receive Nucala® (mepolizumab) if patient has signs/symptoms of parasitic infection, is currently being treated for a parasitic infection, or is having acute bronchospasm and/or asthma attack.

1. Patient Name: _____ Height (inches): _____ Weight (lbs): _____

2. Allergies: _____

3. Diagnosis:

- J45.50 Severe persistent asthma, uncomplicated J45.51 Severe persistent asthma with (acute) exacerbation
 J45.52 Severe persistent asthma with status asthmaticus
 Other ICD-10 Code/description: _____

Provider Attestation for HCP administration:

- Provider attestation that the patient or caregiver are not competent or are physically unable to administer the Nucala product FDA labeled for self-administration
 Patient has experienced severe hypersensitivity reactions (e.g., anaphylaxis, angioedema, bronchospasm, or hypotension) to Nucala within the past 6 months and requires administration and direct monitoring by a healthcare professional
 Patient has a history of uncontrolled disease and ordering provider attests that in their clinical opinion, it is not advisable to try the self-administered formulation of requested drug
 Due to patient's weight, ordering provider attests that in their clinical opinion, it is not advisable to try the self-administered formulation of requested drug
 The location and circumstances for self-administration are not adequate for the potential treatment of anaphylaxis should that arise.

Dose/Frequency:

Nucala® (mepolizumab) 100 mg every four (4) weeks via subcutaneous injection
(Administer as subcutaneous injection only to upper arm, thigh, or abdomen)

Special Orders: _____

Extended post treatment monitoring for any patient new to therapy: monitor patient for one (1) hour after first injection, for 30- minutes after second injection, and then 15-minutes after each injection thereafter.

If adverse drug reaction occurs, utilize the ADVERSE DRUG REACTION GUIDELINES

4. Physician's Signature: _____ / _____ Date: _____
No Stamp Signatures (Dispense as written) (Substitution permitted)

Printed Physician's Name with Credentials: _____ NPI: _____

5. Fax updated supporting clinical MD notes with each order renewal or change in orders
Infusion order forms and Adverse Drug Reaction Guidelines are available at www.palmettoinfusion.com

MRN: _____

DOB: _____

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Guidelines for Prescribing Nucala® (mepolizumab)

(Required documentation with all initial referrals)

Patient Name: _____

Referral Date: _____

- Include signed and completed **Plan of Treatment**. (MD must complete sections 1-5)
(Infusion order forms & Standard Adverse Reactions orders are available at www.palmettoinfusion.com under Agency/MD tab)
- Include patient demographic information and insurance information. (Copy of insurance cards if available)
- Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy. Include any documented use of inhaled or oral corticosteroids, glucocorticoids, theophylline, leukotriene modifiers, short or long acting beta-agonists. Any lab results and/or Pulmonary Function Tests to support diagnosis.**
 - NUCALA® is an interleukin-5 antagonist monoclonal antibody (IgG1 kappa) indicated for: Add-on maintenance treatment of patients with severe asthma aged 12 years and older, and with an eosinophilic phenotype.
- If patient is switching biologic therapies such as Xolair® or Cinqair®, then MD must specify wash-out period prior to starting NUCALA® as specified of _____ weeks.
Last known therapy: _____ and last known date received: _____.
- Other as requested: _____

Pre-Screening:

- Blood Eosinophil Level (Pre-treatment baseline count greater than or equal to 150 cells/mcL)
(Absolute Eosinophil in K/ μ L x 1000 = cells/mcL)

**** Warnings/Precautions: Hypersensitivity reactions:** (e.g., anaphylaxis, angioedema, bronchospasm, hypotension, urticaria, rash) have occurred after administration of NUCALA®. Discontinue NUCALA® in the event of a hypersensitivity reaction. Do not use to treat acute bronchospasm or status asthmaticus. The most common adverse reactions (incidence \geq 5%) include headache, injection site reaction, back pain, and fatigue. **Herpes zoster infections:** have occurred in patients receiving NUCALA®. Consider vaccination if medically appropriate. **Helminth Infections:** Treat patients with pre-existing parasitic infections before therapy. If patients become infected while receiving treatment with NUCALA® and do not respond to anti-helminth treatment, then discontinue NUCALA® until parasitic infection resolves. **Corticosteroids:** Do not discontinue systemic or inhaled corticosteroids abruptly upon initiation of therapy. Decrease corticosteroids gradually, if appropriate. **Pregnancy/Breastfeeding:** Discuss Pregnancy or breastfeeding plans/risks prior to start of therapy. See full prescribing information.

AccuRx Infusion will complete insurance verification and submit all required clinical documentation to the patient’s insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.

Please fax all information to 1-866-872-8920 or call 1-800-809-1265 for assistance.



Checklist for referrals to AccuRX Infusion Center:

Fax referral to 1-866-990-3192

- Patient demographics – address, phone number, SS#, etc.
- Insurance Information – copy of the card(s) if possible
- Plan of Treatment/Orders
- Most recent physician office notes to include tried and failed therapies – all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs
- Any lab results or other diagnostic procedures to support the diagnosis

AccuRx Infusion Center will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility. Our office will notify you if any further information is required. We will review financial responsibility with the patient and refer them to any available Co-pay assistance as required. Thank you for the referral.