

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/>	<input type="checkbox"/> Order Renewal
Patient preferred clinic:			

Cathflo® Activase® (alteplase) Standard Plan of Treatment for Line Care

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight:	LB or KG	Gender: Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NDKA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="text"/>	- Other:
<input type="text"/>	- Other:

REQUESTED DOCUMENTATION:

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INJECTION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INJECTION DATE:
4		FROM PREVIOUS	
5		THERAPY:	
6			

Continue current order until insurance approved

MEDICATION ORDERS:

MEDICATION/DOSE:

- Cathflo® Activase® (alteplase) 2mg intracatheter, x ____ lumen(s).
 Instill alteplase for no blood return, occluded line, or sluggish flush. After 30 minutes of dwell time, reassess catheter patency by aspirating blood return. If catheter function is not restored after 120 minutes of dwell time, then a second dose may be instilled. **Patients weighing less than 30 kg require adjusted dosing.**

Catheter Specific Orders:

- Change dressing and cap(s) as required per protocol. If catheter function is restored, aspirate 4-5 ml of blood in patients greater than or equal to 10 kg or 3 ml in patients less than 10 kg to remove Cathflo® Activase® (alteplase) and residual clot. Flush line with 5-20ml of Sodium Chloride 0.9% IV per line type as required. Then flush with Heparin 100 units/ml IV per line type as required or **Heparin 10 units/ml IV 1-5ml per line type as required (for pediatric patients)**. Notify referring physician if patency is not established and further clinical evaluation is required.

SPECIAL ORDERS:

<input type="text"/>

LINE USE/CARE ORDERS:

- Start PIV/Access CVC
 Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE

<input type="text"/>	<input type="text"/>
Dispense as written/Brand medically necessary	Substitution permitted