Palmetto INFUSION®

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Referral Status:			MRN:
☐ New referral	☐ Order change	☐ Order Renewal	
Patient preferred clinic:			

Cimzia® (certolizumab pegol) Standard Plan of Treatment for Gastroenterology

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches: / Weight: LB or KG	Gender: / Allergies: ☐ See list ☐ NDKA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

- ☐ K50.____ - Crohn's disease (small instestine)
- ☐ K50.1 - Crohn's disease (large intestine)
- ☐ K50.8 - Crohn's disease (small & large intestine)
- ☐ K50.9 - Crohn's disease, unspecified
- ☐ ________ - Other: ____

REQUESTED DOCUMENTATION:

1	Insurance information
2	Most recent History & Physical
3	Full medication list
4	Tried and failed therapies
5	TB screening
6	HBV screening/labs as required by payor

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

IF NO:	IF YES:
PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INJECTION DATE:
	NEXT INJECTION DATE:

IF ORDER CHANGE:

☐ **Continue current order until insurance approved**

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive certolizumab pegol if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

DOSE/FREQUENCY:

☐ Induction: Cimzia® (certolizumab pegol) 400mg at week 0, week 2, week 4, and every 4 weeks thereafter

☐ Maintenance: Cimzia® (certolizumab pegol) 400mg every 4 weeks

Administer as 2 divided subcutaneous injections to separate sites in the abdomen or thigh only

SPECIAL ORDERS:

☐ ____

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment. Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.

☑ Refills x 12 months unless noted otherwise here:

LINE USE/CARE ORDERS:

☑ Start PIV/Access CVC

☑ Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures) DATE

Dispense as written/Brand medically necessary	Substitution permitted	