

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

Cimzia® (certolizumab pegol) Standard Plan of Treatment for Gastroenterology

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
<input type="checkbox"/> See list	<input type="checkbox"/> NDKA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="checkbox"/> K50.____ - Crohn's disease (small intestine)
<input type="checkbox"/> K50.1 - Crohn's disease (large intestine)
<input type="checkbox"/> K50.8 - Crohn's disease (small & large intestine)
<input type="checkbox"/> K50.9 - Crohn's disease, unspecified
<input type="checkbox"/> _____ - Other:

REQUESTED DOCUMENTATION: PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INJECTION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INJECTION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	TB screening	THERAPY:	
6	HBV screening/labs as required by payor		
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive certolizumab pegol if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

DOSE/FREQUENCY:

- Induction: Cimzia® (certolizumab pegol) 400mg at week 0, week 2, week 4, and every 4 weeks thereafter
- Maintenance: Cimzia® (certolizumab pegol) 400mg every 4 weeks

Administer as 2 divided subcutaneous injections to separate sites in the abdomen or thigh only

SPECIAL ORDERS:

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment.
 Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.

Refills x 12 months unless noted otherwise here:

LINE USE/CARE ORDERS: ADVERSE REACTION & ANAPHYLAXIS ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures) DATE

Dispense as written/Brand medically necessary	Substitution permitted	



Palmetto

INFUSION

Checklist for referrals to Palmetto Infusion:

Fax referral to 1.866.872.8920

- Patient demographics – address, phone number, SS#, etc.**
- Insurance Information – copy of the card(s) if possible**
- Plan of Treatment/Orders**
- Most recent physician office notes to include tried and failed therapies – all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs.**
- Any lab results or other diagnostic procedures to support the diagnosis**

Palmetto Infusion will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility.

Our office will notify you if any further information is required.

We will review financial responsibility with the patient and refer them to any available co-pay assistance as required. Palmetto Infusion Call Center 800.809.1265. Thank you for the referral.

www.PalmettoInfusion.com