

Entyvio® (vedolizumab) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: See list NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

K50.0 - Crohn's Disease (small intestine)	K51.0 - Universal Ulcerative (chronic) Pancolitis
K50.1 - Crohn's Disease (large intestine)	K51.8 - Other Ulcerative (chronic) Colitis
K50.8 - Crohn's Disease (small & large intestine)	K51.5 - Left sided Ulcerative (chronic) Colitis
K50.9 - Crohn's Disease	K51.9 - Ulcerative Colitis
- Other:	

REQUESTED DOCUMENTATION:

1 Insurance information	IF NO:	IF YES:
2 Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3 Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4 Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	THERAPY:	Continue current order until insurance approved
6		

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive vedolizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV	Diphenhydramine	25mg	50mg	Other:	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
					Loratadine	10mg				
					Other:					

MEDICATION/DOSE:

☒ Entyvio® (vedolizumab) 300mg per 250ml NS IV to infuse over at least 30 minutes

FREQUENCY:

☐ Induction: Dosed at week 0, week 2, week 6, and every 8 weeks thereafter

☐ Maintenance: Dosed every 8 weeks

☐ Other: _____

SPECIAL/LAB ORDERS:

☐ _____



Refills x 12 months unless noted otherwise here:

LINE USE/CARE ORDERS:

☒ Start PIV/Access CVC

☒ Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted
---	------------------------