

Referral Status:				MRN:	
<input type="checkbox"/>	New referral	<input type="checkbox"/>	Order change	<input type="checkbox"/>	Order Renewal
Patient preferred clinic:					

Infliximab Unspecified Plan of Treatment for Dermatology

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:				
Patient Name:		Address:				
Date of Birth:		City, State, Zip:				
Height in inches:	Weight: LB or KG	Gender:	Allergies:		See list	NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

	L40.5	- Psoriatic Arthritis/Arthropathy
	L40.	- Psoriasis
		- Other:

REQUESTED DOCUMENTATION:

1	Insurance information
2	Most recent History & Physical
3	Full medication list
4	Tried and failed therapies
5	<u>REQUIRED:</u> TB screening for new start
6	HBV screening/labs as required by payor

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION

IF NO:	IF YES:	
PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:	
	NEXT INFUSION DATE:	
	IF ORDER CHANGE:	
		Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive infliximab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

Premedication with antihistamines, acetaminophen, and/or corticosteroids may be considered to prevent infusion-related reactions.

IV	Diphenhydramine	25mg	50mg		PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
SPECIFIC MEDICATION:						Cetirizine	10mg			
	Remicade	<u>Any infliximab</u>				Loratadine	10mg			
	Avsola	biosimilar may be				Other:				

SPECIFIC MEDICATION:

	Remicade
	Avsola
	Inflectra
	Renflexis

DOSE:

<input type="checkbox"/>	5mg/kg diluted in NS infused IV per step protocol over 2 hours
<input type="checkbox"/>	Other: _____
<input type="checkbox"/>	May utilize expedited infusion per step protocol to run over 1 hour as tolerated

FREQUENCY:

<input type="checkbox"/>	Induction to be completed at week 0, week 2, and week 6, and then every 8 weeks thereafter
<input type="checkbox"/>	Maintenance every 8 weeks
<input type="checkbox"/>	Infuse every _____ weeks

SPECIAL/LAB ORDERS:

☐ _____

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment.

Prescriber to monitor patient for symptoms of HBV and TB infection and reactivation as clinically appropriate.

☒ Refills x 12 months unless noted otherwise here:

LINE USE/CARE ORDERS:

- ☒ Start PIV/Access CVC
- ☒ Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.

**PRESCRIBER INFORMATION:**

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

		DATE:
Dispense as written/Brand medically necessary		Substitution permitted