

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
Patient preferred clinic:	<input type="checkbox"/> Order change

Ocrevus® (ocrelizumab) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
	<input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="checkbox"/> G35 - Relapsing Remitting Multiple Sclerosis
<input type="checkbox"/> G35 - Primary Progressive Multiple Sclerosis
<input type="checkbox"/> - Other:

REQUESTED DOCUMENTATION: PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	REQUIRED: Hepatitis B Panel for new start patients	THERAPY:	
6	Quantitative Serum Immunoglobulin screening		Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive ocrelizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

*Per FDA labeling, Acetaminophen PO, Diphenhydramine IVP, and Methylprednisolone IVP is suggested prior to infusion

IV	Diphenhydramine	25mg	50mg	PO	Acetaminophen	325mg	500mg	650mg	1000mg	
	Methylprednisolone	40mg	125mg		Other:	Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
					Cetirizine	10mg				
					Loratadine	10mg				
					Other:					

DOSE AND FREQUENCY: PRODUCT:

Induction: Ocrevus® 300mg IV in 250ml NS to be infused over 2.5 hours or longer per step protocol at week 0 and 2 weeks

Maintenance: Ocrevus® 600mg IV in 500ml NS every 6 months

SPECIAL/LAB ORDERS:

MAINTENANCE INFUSION TIME:

Infuse maintenance dose over 2 hours per step protocol


Infuse maintenance dose over 3.5 - 4 hours per step protocol

Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.

Refills x 12 months unless noted otherwise here:

***Maintenance dosing is scheduled 6 months from initial 0-week dosing.**

LINE USE/CARE ORDERS: ADVERSE REACTION & ANAPHYLAXIS ORDERS:

<input checked="" type="checkbox"/> Start PIV/Access CVC <input checked="" type="checkbox"/> Flush device per facility standard flushing procedure	Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.	
---	---	---

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures) DATE

Dispense as written/Brand medically necessary	Substitution permitted