

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

ONPATTRO™ (patisiran) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
<input type="checkbox"/> See list	<input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

E85.1 - Neuropathic Heredofamilial amyloidosis
- Other:

REQUESTED DOCUMENTATION:

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	Labs/Tests supporting primary diagnosis (serum TTR, PND Scores, FAP stage, or modified Neuropathy Impairment Scores)	THERAPY:	
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive ONPATTRO™ if demonstrating signs and symptoms suggestive of vitamin A deficiency.

PREMEDICATION: To be administered 60 minutes prior to infusion as selected.

*FDA labeling suggests that all patients are premedicated with IV corticosteroid, acetaminophen 500mg PO, and both H1 and H2 antihistamine blocker IV 60 minutes prior to infusion as per selected by referring physician below.

IV	Diphenhydramine	25mg	50mg	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg		Other:	Famotidine	20mg	40mg	
	Famotidine	20mg	40mg			Diphenhydramine	25mg	50mg	
	Dexamethasone	10mg	Other:			Fexofenadine	60mg	180mg	
	Other:					Cetirizine	10mg		
						Loratadine	10mg		
				Other:					

MEDICATION:

ONPATTRO™ (patisiran) in NS for a total volume of 200 ml IV via pump as per step protocol. **Utilizing infusion set and line that are DEHP-free.**

DOSE /FREQUENCY:

<input type="checkbox"/>	<100kg: 0.3mg/kg IV every 3 weeks
<input type="checkbox"/>	>100kg: 30mg IV every 3 weeks
<input type="checkbox"/>	Other: _____

SPECIAL/LAB ORDERS:

<input type="checkbox"/>	_____
--------------------------	-------

*If dose is received within 3 days of missed dose, then continue dosing according to original schedule. If greater than 3 days after missed dose, then continue dosing every 3 weeks thereafter.

<input checked="" type="checkbox"/>	Refills x 12 months unless noted otherwise here:
-------------------------------------	--

LINE USE/CARE ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted