Palmetto INFUSION®

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Referral Status:		MRN:
☐ New referral	☐ Order change	☐ Order Renewal
Patient preferred clinic:		

Orencia® (abatacept) Plan of Treatment for Rheumatology

PATIENT DEMOGRAPHICS:

Date of Referral:			Patient's Phone:		
Patient Name:			Address:		
Date of Birth:			City, State, Zip:		
Height in inches:	Weight: LB or KG		Gender:	Allergies: ☐	See list ☐ NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

☐ M05._____ - Rheumatoid Arthritis with Rheumatoid factor

☐ M06._____ - Rheumatoid Arthritis without Rheumatoid factor

☐ _____ - Other:

REQUESTED DOCUMENTATION:		PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?	
1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:
3	Full medication list		NEXT INFUSION DATE:
4	Tried and failed therapies		**IF ORDER CHANGE:**
5	TB screening results		☐ **Continue current order until insurance approved**
6	HBV screening/labs as required by payor		

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive abatacept if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening diagnosis of COPD or respiratory status, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV	Diphenhydramine	☐ 25mg	☐ 50mg						
	Methylprednisolone	☐ 40mg	☐ 125mg	☐ Other:					
	Famotidine	☐ 20mg	☐ 40 mg						
	Other:	☐							
PO	Acetaminophen	☐ 325mg	☐ 500mg	☐ 650mg	☐ 1000mg				
	Famotidine	☐ 20mg	☐ 40mg						
	Diphenhydramine	☐ 25mg	☐ 50mg						
	Fexofenadine	☐ 60mg	☐ 180mg						
	Cetirizine	☐ 10mg							
	Loratadine	☐ 10mg							
	Other:	☐							

MEDICATION:

☑ Orencia® (abatacept) dosage per 100 ml NS to infuse over at least 30 minutes.

DOSE:

☐ Dosage based on the following guidelines from the FDA package labeling.

Patient Weight	Dose	250 mg vials
<60 kg	500 mg	2
60 to 100kg	750 mg	3
>100 kg	1000 mg	4

☐ Flat Dose: _____mg

FREQUENCY:

☐ Induction: To be given at 0 week, 2 week, and 4 weeks

☐ Maintenance: Every 4 weeks

☐ Other:_____

SPECIAL/LAB ORDERS:

☐ _____

_____

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment. Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.

☑ Refills x 12 months unless noted otherwise here:

LINE USE/CARE ORDERS:

☑ Start PIV/Access CVC

☑ Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)		DATE:
Dispense as written/Brand medically necessary	Substitution permitted	