

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

Skyrizi® (risankizumab-rzaa) Standard Plan of Treatment for Gastroenterology

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

K50.0 - Crohn's Disease (small intestine)	K50.1 - Crohn's Disease (large intestine)
K50.8 - Crohn's Disease (small & large intestine)	K50.9 - Crohn's Disease
- Other: _____	

REQUESTED DOCUMENTATION:

1	Insurance information
2	Most recent History & Physical
3	Full medication list
4	Tried and failed therapies
5	REQUIRED: TB screening for new start patients
6	Baseline LFTs and bilirubin level

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

IF NO:	IF YES:
PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:
	NEXT INFUSION DATE:
	IF ORDER CHANGE:
	Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive risankizumab-rzaa if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new-onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV	Diphenhydramine	25mg	50mg	Other:	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
						Loratadine	10mg			
						Other:				

MEDICATION/DOSE:

☒ Skyrizi® (risankizumab-rzaa) 600mg/10ml in 100ml-500ml of D5W given IV over at least 1 hour

FREQUENCY:

☐ Week 0, week 4, and week 8
☐ Other: _____

SPECIAL/OTHER LAB ORDERS:

LINE USE/CARE ORDERS:

- ☒ Start PIV/Access CVC
☒ Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

_____ Dispense as written/Brand medically necessary	_____ Substitution permitted	_____
--	---------------------------------	-------