

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

Skyrizi® (risankizumab-rzaa) Standard Plan of Treatment for Gastroenterology

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

K50.0 - Crohn's Disease (small intestine)	K50.1 - Crohn's Disease (large intestine)
K50.8 - Crohn's Disease (small & large intestine)	K50.9 - Crohn's Disease
- Other: _____	

REQUESTED DOCUMENTATION:

REQUESTED DOCUMENTATION:	PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?
1 Insurance information	IF NO: _____ IF YES: _____
2 Most recent History & Physical	PLEASE STATE LAST INFUSION DATE: _____
3 Full medication list	REQUIRED WASHOUT FROM PREVIOUS THERAPY: _____ NEXT INFUSION DATE: _____
4 Tried and failed therapies	IF ORDER CHANGE: <input type="checkbox"/> Continue current order until insurance approved
5 Baseline liver enzymes	
6 Baseline bilirubin levels	

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive risankizumab-rzaa if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new-onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV	Diphenhydramine	25mg	50mg	Other: _____	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other: _____					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
				Loratadine	10mg					
				Other: _____						

MEDICATION/DOSE:

Skyrizi® (risankizumab-rzaa) 600mg/10ml in 100ml-500ml of D5W given IV over at least 1 hour

SPECIAL/OTHER LAB ORDERS:

FREQUENCY:

Week 0, week 4, and week 8

Other: _____

LINE USE/CARE ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted



Palmetto

INFUSION

Checklist for referrals to Palmetto Infusion:

Fax referral to 1.866.872.8920

- Patient demographics – address, phone number, SS#, etc.**
- Insurance Information – copy of the card(s) if possible**
- Plan of Treatment/Orders**
- Most recent physician office notes to include tried and failed therapies – all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs.**
- Any lab results or other diagnostic procedures to support the diagnosis**

Palmetto Infusion will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility.

Our office will notify you if any further information is required.

We will review financial responsibility with the patient and refer them to any available co-pay assistance as required. Palmetto Infusion Call Center 800.809.1265. Thank you for the referral.

www.PalmettoInfusion.com