

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

## Soliris® (eculizumab) Standard Plan of Treatment for Pediatric aHUS

### PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
	See list
	NKDA

### DIAGNOSIS: (PLEASE COMPLETE 2<sup>ND</sup> AND 3<sup>RD</sup> DIGITS TO COMPLETE ICD 10 FOR BILLING)

D59.3 - Atypical Hemolytic Uremic Syndrome (aHUS)	D59.4 - Other non autoimmune hemolytic anemias (including microangiopathic hemolytic anemia)
D58.8 - Other specified hereditary hemolytic anemias	
D59.8 - Other acquired hemolytic anemias	
- Other:	

### REQUESTED DOCUMENTATION:

### PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:
3	Full medication list		NEXT INFUSION DATE:
4	Tried and failed therapies		<b>IF ORDER CHANGE:</b>
5	Documented meningococcal vaccine administration (Both MenACWY AND MenB)		<b>Continue current order until insurance approved</b>

### MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive eculizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, presents with any symptoms of meningococcal infections, and/or surgery.

### PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

\*FDA labeling does not suggest any premedication prior to infusion

IV	Diphenhydramine	25mg	50mg		PO	Acetaminophen	325mg	500mg	160mg/5ml	mls
	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg	12.5mg/5ml:	mls
	Other:					Loratadine	10mg			

### MEDICATION:

Soliris® (eculizumab) IV given over 35 minutes diluted in NS according to FDA labeling suggestions.

**If the infusion is slowed, the total infusion time should not exceed 2 hours.**

\*Follow each infusion with a 1 hour post infusion monitoring\*

### FREQUENCY/DOSE:

Patient body weight	Induction	Maintenance
≥ 40kg	900 mg weekly x 4 doses	1200 mg at week 5; then 1200 mg every 2 weeks thereafter
30kg to less than 40kg	600 mg weekly x 2 doses	900 mg at week 3; then 900 mg every 2 weeks thereafter
20kg to less than 30kg	600 mg weekly x 2 doses	600 mg at week 3; then 600 mg every 2 weeks thereafter
10kg to less than 20kg	600 mg weekly x 1 dose	300 mg at week 2; then 300 mg every 2 weeks thereafter
5kg to less than 10kg	300 mg weekly x 1 dose	300 mg at week 2; then 300 mg every 3 weeks thereafter

**Prescriber must be enrolled in the Soliris (REMS) program, at 18887654747 or at [www.solirisrems.com](http://www.solirisrems.com).**

Refills x 12 months unless noted otherwise here:

### LINE USE/CARE ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure

### ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



### PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

### PRESCRIBER SIGNATURE: (No stamp signatures)

### DATE

Dispense as written/Brand medically necessary	Substitution permitted