

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Preferred clinic:	DAYS IN AIC: DAYS IN HOME:

## Tzield® (teplizumab-mzww) Plan of Treatment

### PATIENT DEMOGRAPHICS:

Patient Name:	Patient's Phone:
Date of Birth:	Address:
Allergies:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

### DIAGNOSIS: (PLEASE COMPLETE 2<sup>ND</sup> AND 3<sup>RD</sup> DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="checkbox"/> E10.9 - Type 1 diabetes mellitus without complications	<input type="checkbox"/> -Other:
<input type="checkbox"/> E10.8 - Type 1 diabetes mellitus with unspecified complications	

### REQUESTED DOCUMENTATION:

REQUESTED DOCUMENTATION:	HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE? Yes or No
1 Insurance information	PLEASE STATE
2 H&P including tried and failed therapies	REQUIRED WASHOUT
3 Full medication list	FROM PREVIOUS
4 <b>Required: Recent CBC with diff and LFTs</b>	OTHER:

**Continue current order until insurance approved**

### HOME SUPPLY ORDER: \*Applicable to all home infusion patients subject to insurance approval

All supplies for vascular access line care, dressing kit, drug administration, adverse reaction kit, Infusion pump, IV pole, pole clamp etc. will be

### MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive teplizumab-mzww if receiving antibiotics for active infectious process, active fever and/or suspected infection or surgery. If patient experiences severe cytokine release syndrome, consider temporarily pausing therapy for 1-2 days and review prescribing information before continuing therapy.

<input checked="" type="checkbox"/>	<b>Home Anaphylaxis Kit:</b> Dispense and administer for mild and severe reaction. Applicable to all home infusion patients.
2 - Epinephrine 1 mg/ml 1 ml 2- Diphenhydramine 50 mg/ml 2 ml vial Syringes, needles and 0.9% Normal Saline Flushes 10 mls to administer Complete Home Infusion Physician Standing Order for complete Home Infusion, Nursing, and Adverse Reaction Orders to be utilized in the event of an adverse reaction/anaphylaxis. Epinephrine administered IM per weight based dosing guide and Benadryl 25-50mg IVP to be administered by clinician in the home.	

**\*Per FDA labeling: Premedication of Acetaminophen or NSAID and antihistamine, and/or an antiemetic is suggested for at least the first 5 days of the 14-day treatment course**

**\*Any selected premedication will only be given with infusions 1 through 5 unless otherwise designated here:**

<b>IV</b>	Diphenhydramine	25mg	50mg		<b>PO</b>	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Ondansetron	4mg	8mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
				Loratadine	10mg					
				Other:						

### DOSAGE:

<input checked="" type="checkbox"/>	Treatment Day	Dosage
	Day 1	65 mcg/m2
	Day 2	125 mcg/m2
	Day 3	250 mcg/m2
	Day 4	500 mcg/m2
	Day 5-14	1030 mcg/m2

### LAB ORDERS:

<input type="checkbox"/>	Draw CBC with diff & LFT on infusion days 5 and 8
<input type="checkbox"/>	Other: _____

### ADMINISTRATION:

Tzield® 100 mcg/mL diluted in 25mL NS administered as an IV infusion over a minimum of 30-minutes followed by a 25mL NS flush. Administer once daily for 14 consecutive days.

Follow first 5 infusions with a 1-hour post infusion observation.

### Physician responsible for all follow up lab monitoring

Per FDA labeling, with AST/ALT > 5x ULN, bilirubin > 3x ULN, or Lymphocyte count <500/mcL lasting one week or longer, patient may be ineligible to receive Tzield®

### SITE OF CARE:

<input type="checkbox"/> AIC	<input type="checkbox"/> AIC and Home with a nurse
------------------------------	----------------------------------------------------

*\*If site of care not indicated, PIS will coordinate with patient*

### SPECIAL/OTHER LAB ORDERS:

<input type="checkbox"/>	_____
<input type="checkbox"/>	_____

### LINE USE/CARE ORDERS:

<input checked="" type="checkbox"/> Start PIV/Access CVC <input checked="" type="checkbox"/> Flush PIV/Access per PIV/PICC/CVC protocol.	<b>Dispense and Administer as Prescribed</b>
---------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------

### PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

### PRESCRIBER SIGNATURE: (No stamp signatures)

### DATE:

Dispense as written/Brand medically necessary	Substitution permitted