

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

Ocrevus® (ocrelizumab) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="checkbox"/> G35 - Relapsing Remitting Multiple Sclerosis
<input type="checkbox"/> G35 - Primary Progressive Multiple Sclerosis
<input type="checkbox"/> - Other:

REQUESTED DOCUMENTATION:

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	Quantitative Serum Immunoglobulin screening	THERAPY:	
6	REQUIRED: HBsAg, anti-HBc, and anti-HBs		
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive ocrelizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

*Per FDA labeling, Acetaminophen PO, Diphenhydramine IVP, and Methylprednisolone IVP is suggested prior to infusion

IV	Diphenhydramine	25mg	50mg		PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
					Cetirizine	10mg				
					Loratadine	10mg				
					Other:					

MEDICATION/FREQUENCY:

Induction: Ocrevus® 300mg IV in 250ml NS to be infused over 2.5 hours or longer per step protocol at week 0 and 2 weeks

Maintenance: Ocrevus® 600mg IV in 500ml NS every 6 months

Follow each infusion with a (1) one-hour post observation period.

MAINTENANCE INFUSION TIME:

Infuse maintenance dose over 2 hours per step protocol

Infuse maintenance dose over 3.5 - 4 hours per step protocol

Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.



Refills x 12 months unless noted otherwise here:

*Maintenance dosing is scheduled 6 months from initial 0-week dosing.

LINE USE/CARE ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE

Dispense as written/Brand medically necessary	Substitution permitted



Checklist for referrals to Palmetto Infusion: Fax referral to 1.866.872.8920

- Patient demographics - address, phone number, SS#, etc.**
- Insurance information - copy of the card(s) if possible**
- Plan of Treatment/Orders**
- Most recent physician office notes to include tried and failed therapies - all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs.**
- Any lab results or other diagnostic procedures to support the diagnosis.**

Palmetto Infusion will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility.

Our office will notify you if any further information is required.

We will review financial responsibility with patient and refer them to any available co-pay assistance as required. Palmetto Infusion Call Center 800.809.1265. Thank you for your referral.

www.PalmettoInfusion.com