Referral Status:			MRN:
☐ New referral	☐ Order change	☐ Order Renewal	
Patient preferred clinic:			

Simponi ARIA® (golimumab) Standard Plan of Treatment for Rheumatology

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches: / Weight: LB or KG	Gender: / Allergies: ☐ See list ☐ NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

☐	M05._____ - Rheumatoid Arthritis with Rheumatoid factor	☐	M06._____ - Rheumatoid Arthritis without Rheumatoid factor
☐	L40.5_____ - Psoriatic Arthropathy	☐	M45._____ - Ankylosing Spondylitis
☐	________- Other:________________________________		

REQUESTED DOCUMENTATION:

1	Insurance information
2	Most recent History & Physical
3	Full medication list
4	Tried and failed therapies
5	**REQUIRED:** TB screening for new start
6	HBV screening/labs as required by payor

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

IF NO:	IF YES:
PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:
	NEXT INFUSION DATE:
	IF ORDER CHANGE:
	☐ **Continue current order until insurance approved**

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive golimumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new onset or deterioration neurological changes, and/or surgery

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV				
Diphenhydramine	☐ 25mg	☐ 50mg		
Methylprednisolone	☐ 40mg	☐ 125mg	☐ Other:	
Famotidine	☐ 20mg	☐ 40 mg		
Other:	☐			

PO				
Acetaminophen	☐ 325mg	☐ 500mg	☐ 650mg	☐ 1000mg
Famotidine	☐ 20mg	☐ 40mg		
Diphenhydramine	☐ 25mg	☐ 50mg		
Fexofenadine	☐ 60mg	☐ 180mg		
Cetirizine	☐ 10mg			
Loratadine	☐ 10mg			
Other:	☐			

MEDICATION/DOSE:

☑ Simponi ARIA® (golimumab) 2 mg/kg per 100 ml NS given IV to infuse over at least 30 minutes

FREQUENCY:

☐ Induction: Given at 0 week and 4 weeks, and then every 8 weeks thereafter

☐ Maintenance: Given every 8 weeks

☐ Other: ______________________________________

SPECIAL/OTHER LAB ORDERS:

☐ ____

____

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment. Prescriber to monitor patient for symptoms of HBV and TB infection and reactivation as clinically appropriate.

☑ Refills x 12 months unless noted otherwise here:



LINE USE/CARE ORDERS:

☑ Start PIV/Access CVC

☑ Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

		DATE:
Dispense as written/Brand medically necessary	Substitution permitted	

Checklist for referrals to Palmetto Infusion: Fax referral to 1.866.872.8920

- [ ] **Patient demographics – address, phone number, SS#, etc.**
- [ ] **Insurance information – copy of the card(s) if possible**
- [ ] **Plan of Treatment/Orders**
- [ ] **Most recent physician office notes to include tried and failed therapies – all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs.**
- [ ] **Any lab results or other diagnostic procedures to support the diagnosis.**

Palmetto Infusion will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility.

Our office will notify you if any further information is required.

We will review financial responsibility with patient and refer them to any available co-pay assistance as required. Palmetto Infusion Call Center 800.809.1265. Thank you for your referral.

www.PalmettoInfusion.com