

| | |
|--|---------------------------------------|
| Referral Status: | MRN: |
| <input type="checkbox"/> New referral | <input type="checkbox"/> Order change |
| <input type="checkbox"/> Order Renewal | |
| Patient preferred clinic: | |

Cimzia® (certolizumab pegol) Standard Plan of Treatment for Rheumatology & Dermatology

PATIENT DEMOGRAPHICS:

| | |
|-------------------|-------------------|
| Date of Referral: | Patient's Phone: |
| Patient Name: | Address: |
| Date of Birth: | City, State, Zip: |
| Height in inches: | Weight: LB or KG |
| Gender: | Allergies: |
| | See list |
| | NKDA |

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

| | |
|---|--|
| M05. ___ - Rheumatoid Arthritis with rheumatoid factor | M45.A ___ - Non-radiographic axial spondyloarthritis |
| M06. ___ - Rheumatoid Arthritis without rheumatoid factor | L40.0 - Psoriasis vulgaris |
| M45. ___ - Ankylosing Spondylitis | L40.5 ___ - Arthropathic psoriasis |
| M46.8 ___ - Other specified inflammatory spondylopathies | L40.9 - Psoriasis, unspecified |
| ___ - Other: | |

REQUESTED DOCUMENTATION:

| | |
|---|--|
| 1 | Insurance information |
| 2 | Most recent History & Physical |
| 3 | Full medication list |
| 4 | Tried and failed therapies |
| 5 | REQUIRED: TB screening for new start patients |
| 6 | HBV screening/labs as required by payor |

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

| | |
|------------------|--|
| IF NO: | IF YES: |
| PLEASE STATE | LAST INJECTION DATE: |
| REQUIRED WASHOUT | NEXT INJECTION DATE: |
| FROM PREVIOUS | |
| THERAPY: | |
| | IF ORDER CHANGE: |
| | Continue current order until insurance approved |

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive certolizumab pegol if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new-onset or deterioration neurological changes, and/or surgery.

DOSE/FREQUENCY:

Induction: Cimzia® (certolizumab pegol) 400mg at week 0, week 2, week 4, and every 4 weeks thereafter

Maintenance: Cimzia® (certolizumab pegol) 200mg every 2 weeks

Maintenance: Cimzia® (certolizumab pegol) 400mg every 4 weeks

Administer as 2 divided subcutaneous injections to separate sites in the abdomen or thigh only

SPECIAL ORDERS:

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment.
 Prescriber to monitor patient for symptoms of HBV and TB infection and reactivation as clinically appropriate.

Refills x 12 months unless noted otherwise here:

NURSING ORDERS:

Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated.

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

| | |
|-------------------|--------|
| PROVIDER NAME: | PHONE: |
| ADDRESS: | FAX: |
| CITY, STATE, ZIP: | NPI: |

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE

| | | |
|---|------------------------|--|
| | | |
| Dispense as written/Brand medically necessary | Substitution permitted | |