

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

### Crysvita® (burosumab-twza) Pediatric Standard Plan of Treatment

**PATIENT DEMOGRAPHICS:**

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

**DIAGNOSIS: (PLEASE COMPLETE 2<sup>ND</sup> AND 3<sup>RD</sup> DIGITS TO COMPLETE ICD 10 FOR BILLING)**

E83.31 - Familial Hypophosphatemia	E83.39 - Other disorders of phosphorus metabolism
- Other: _____	

**REQUESTED DOCUMENTATION: PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?**

1 Insurance information	IF NO:	IF YES:
2 Most recent History & Physical	PLEASE STATE	LAST INJECTION DATE:
3 Full medication list	REQUIRED WASHOUT	NEXT INJECTION DATE:
4 Tried and failed therapies	FROM PREVIOUS	<b>IF ORDER CHANGE:</b>
5 Fasting serum Phosphorus level *required*	THERAPY:	
6 <b>NOTE: Discontinuation of oral phosphate and Vit D analogs 1 week prior to initiation</b>		<b>Continue current order until insurance approved</b>

**MEDICATION ORDERS:**

**NOTE:** Patient may be ineligible to receive Crysvita with elevated phosphorus serum levels. Concomitant use of burosumab-twza with oral phosphate and/or active vitamin D analogs (e.g. calcitriol, paricalcitol, doxercalciferol, calcifediol) due to the risk of hyperphosphatemia.

**MEDICATION:**

Crysvita® (burosumab-twza)  
 (Administered as subcutaneous injection to upper arm, upper thigh, buttocks, or abdomen. Do not give more than 1.5ml per injection site.)

**DOSE:**

Weight ≤ 10kg: 1mg/kg (rounded to the nearest 1mg)  
 Weight ≥ 10kg: 0.8mg/kg up to 90mg (recommended starting dose with a maximum recommended dose of 90mg)  
 Other: \_\_\_\_\_

All doses will be rounded to the nearest 10mg for patients with body weight > 10kg.

**FREQUENCY:**

Every 2 weeks  
 Other: \_\_\_\_\_

**LAB PARAMETERS: (Physician responsible for all follow up lab monitoring)**

- Recommendations:
- Therapy Initiation: Draw fasting serum phosphorus every 4 weeks for first 3 month of treatment
  - Hold medication if serum phosphorus is above 5 mg/dL, redraw levels in 4 weeks, reassess and restart dosing according to package labeling.

**SPECIAL/OTHER ORDERS:**

Hold medication dose if fasting serum phosphorus is greater than \_\_\_\_\_ and call prescriber.

Refills x 12 months unless noted otherwise here:

**LINE USE/CARE ORDERS: ADVERSE REACTION & ANAPHYLAXIS ORDERS:**

Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



**PRESCRIBER INFORMATION:**

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

**PRESCRIBER SIGNATURE: (No stamp signatures) DATE**

_____	_____
Dispense as written/Brand medically necessary	Substitution permitted