

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

Ocrevus® (ocrelizumab) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="checkbox"/> G35 - Relapsing Remitting Multiple Sclerosis
<input type="checkbox"/> G35 - Primary Progressive Multiple Sclerosis
<input type="checkbox"/> - Other:

REQUESTED DOCUMENTATION:

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	Quantitative Serum Immunoglobulin screening	THERAPY:	
6	REQUIRED: HBsAg, anti-HBc, and anti-HBs		
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive ocrelizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

*Per FDA labeling, Acetaminophen PO, Diphenhydramine IVP, and Methylprednisolone IVP is suggested prior to infusion

IV	Diphenhydramine	25mg	50mg		PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
					Cetirizine	10mg				
					Loratadine	10mg				
					Other:					

MEDICATION/FREQUENCY:

Induction: Ocrevus® 300mg IV in 250ml NS to be infused over 2.5 hours or longer per step protocol at week 0 and 2 weeks

Maintenance: Ocrevus® 600mg IV in 500ml NS every 6 months

Follow each infusion with a (1) one-hour post observation period.

MAINTENANCE INFUSION TIME:

Infuse maintenance dose over 2 hours per step protocol

Infuse maintenance dose over 3.5 - 4 hours per step protocol

Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.

Refills x 12 months unless noted otherwise here:

*Maintenance dosing is scheduled 6 months from initial 0-week dosing.

NURSING ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure
- Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE

Dispense as written/Brand medically necessary	Substitution permitted