Palmetto INFUSION®

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Referral Status:			MRN:
☐ New referral	☐ Order change	☐ Order Renewal	
Patient preferred clinic:			

ONPATTRO™ (patisiran) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: ☐ See list ☐ NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

☐	E85.1 - Neuropathic Heredofamilial amyloidosis
☐	_____ - Other:_____

REQUESTED DOCUMENTATION:		PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?	
1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:
3	Full medication list		NEXT INFUSION DATE:
4	Tried and failed therapies		**IF ORDER CHANGE:**
5	Labs/Tests supporting primary diagnosis (serum TTR, PND Scores, FAP stage, or modified Neuropathy Impairment Scores)		☐ **Continue current order until insurance approved**

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive ONPATTRO™ if demonstrating signs and symptoms suggestive of **vitamin A deficiency.**

PREMEDICATION: To be administered 60 minutes prior to infusion as selected.

*FDA labeling suggests that all patients are premedicated with IV corticosteroid, acetaminophen 500mg PO, and both H1 and H2 antihistamine blocker IV 60 minutes prior to Infusion as per selected by referring physician below.

IV	Diphenhydramine	☐	25mg	☐	50mg				
	Methylprednisolone	☐	40mg	☐	125mg	☐	Other:		
	Famotidine	☐	20mg	☐	40mg				
	Dexamethasone	☐	10mg	☐	Other:				
	Other:	☐							

PO	Acetaminophen	☐	325mg	☐	500mg	☐	650mg	☐	1000mg
	Famotidine	☐	20mg	☐	40mg				
	Diphenhydramine	☐	25mg	☐	50mg				
	Fexofenadine	☐	60mg	☐	180mg				
	Cetirizine	☐	10mg						
	Loratadine	☐	10mg						
	Other:	☐							

MEDICATION:

☑ ONPATTRO™ (patisiran) in NS for a total volume of 200 ml IV via pump as per step protocol. Infuse over approximately 80 minutes.
Utilizing infusion set and line that are DEHP-free.

SPECIAL/LAB ORDERS:

☐ _____
_____

DOSE /FREQUENCY:

☐ < 100kg: 0.3mg/kg IV once every 3 weeks
☐ ≥ 100kg: 30mg IV once every 3 weeks
☐ Other:_____

***If dose is received within 3 days of missed dose, then continue dosing according to original schedule. If greater than 3 days after missed dose, then continue dosing every 3 weeks thereafter.**

☑ Refills x 12 months unless noted otherwise here:

LINE USE/CARE ORDERS:

☑ Start PIV/Access CVC
☑ Flush device per facility standard flushing procedure
☑ Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)		DATE:
Dispense as written/Brand medically necessary	Substitution permitted	