

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

Orbactiv® (oritavancin) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: <input type="checkbox"/> See list <input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

<input type="text"/>	- Other:
<input type="text"/>	- Other:

REQUESTED DOCUMENTATION:

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:	LAST INFUSION DATE:
3	Full medication list		NEXT INFUSION DATE:
4	Tried and failed therapies		IF ORDER CHANGE:
5			<input type="checkbox"/> Continue current order until insurance approved
6			

MEDICATION ORDERS:

NOTE: Prescribing Orbactiv® in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria.

Dilution: Use ONLY 5% dextrose in sterile water (D5W) for dilution. **Do NOT use Normal Saline for dilution or flushing of IV line** as it is incompatible with Orbactiv® and may cause precipitation of the drug. Therefore, other intravenous substances, additives, or other medications mixed in normal saline should **NOT** be added to Orbactiv® vials or infused simultaneously through the same IV line or through a common intravenous port. If the same intravenous line is used for sequential infusion of additional medications, the line should be flushed before and after infusion with D5W.

MEDICATION:

- Orbactiv® (oritavancin) in 1000mL of D5W IV to infuse over 3 hours
- Flush with D5W before and after infusion **NS flushes are not compatible**

DOSE/FREQUENCY:

<input type="checkbox"/>	1200mg as one time dose
<input type="checkbox"/>	Other: _____

SPECIAL/LAB ORDERS:

<input type="text"/>	_____
<input type="text"/>	_____

Follow infusion with a 30 minute post observation.



Refills:

NURSING ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure
- Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

<input type="text"/>	<input type="text"/>	<input type="text"/>
Dispense as written/Brand medically necessary	Substitution permitted	



Checklist for referrals to Palmetto Infusion: Fax referral to 1.800.521.9640

- Patient demographics - address, phone number, SS#, etc.**
- Insurance information - copy of the card(s) if possible**
- Plan of Treatment/Orders**
- Most recent physician office notes to include tried and failed therapies - all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs.**
- Any lab results or other diagnostic procedures to support the diagnosis.**

Palmetto Infusion will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility.

Our office will notify you if any further information is required.

We will review financial responsibility with patient and refer them to any available co-pay assistance as required. Palmetto Infusion Call Center 800.809.1265. Thank you for your referral.

www.PalmettoInfusion.com