

Referral Status:		MRN:	
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change	<input type="checkbox"/> Order Renewal	
Patient preferred clinic:			

Tysabri® (natalizumab) Standard Plan of Treatment

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:				
Patient Name:	Address:				
Date of Birth:	City, State, Zip:				
Height in inches:	Weight: LB or KG	Gender:	Allergies:	<input type="checkbox"/> See list	<input type="checkbox"/> NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

G35 - Relapsing Multiple Sclerosis	K50.9 - Crohn's Disease
K50.8 - Crohn's Disease (small & large intestine)	K50.0 - Crohn's Disease (small intestine)
K50.1 - Crohn's Disease (large intestine)	- Other:

REQUESTED DOCUMENTATION:

1 Insurance information	IF NO:	IF YES:
2 Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3 Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4 Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5 Anti-JCV antibodies as required by payer or REMS program	THERAPY:	
		Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive natalizumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new-onset or deterioration neurological changes, and/or surgery

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

*FDA labeling suggests that patients who have stopped treatment for an extended period are at higher risk for hypersensitivity reactions. MD should evaluate premedication and consider antibody testing prior to restart of therapy.

IV	Diphenhydramine	25mg	50mg		PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg		
	Famotidine	20mg	40mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
				Loratadine	10mg					
				Other:						

MEDICATION:

Tysabri® (natalizumab) 300 mg in 100 ml NS IV to infuse over at least 1- hour.

Follow first 12 infusions with a one hour post infusion observation.

FREQUENCY:

Dosing every 4 weeks, no less than every 28 days.
 Other: _____

LAB ORDERS:

Draw JCV antibody test every 6 months
 Draw JCV antibody test every _____

SPECIAL/OTHER LAB ORDERS:

Prescriber to monitor patient for symptoms of PML as clinically appropriate.

*Prior to each infusion: ensure that the patient has a current Notice of Patient Authorization on file to receive Tysabri® (natalizumab) for their diagnosis and complete/submit Pre-infusion Patient Checklist within 24 hours to Biogen Idec

Refills x 12 months unless noted otherwise here:

CARE ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure
- Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted