

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

### Rituximab Unspecified Plan of Treatment for GPA/MPA

#### PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
	<input type="checkbox"/> See list <input type="checkbox"/> NKDA

#### DIAGNOSIS: (PLEASE COMPLETE 2<sup>ND</sup> AND 3<sup>RD</sup> DIGITS TO COMPLETE ICD 10 FOR BILLING)

M31.30 - Granulomatosis with Polyangiitis (GPA/Wegener's Granulomatosis)
M31.7 - Microscopic Polyangiitis (MPA)
- Other:

#### REQUESTED DOCUMENTATION:

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Most recent labs including CBC with diff	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Full medication list / Tried and failed therapies	FROM PREVIOUS	
5	<b>REQUIRED: HBsAg, anti-HBc for new start patients</b>	THERAPY:	
			<b>IF ORDER CHANGE:</b>
			<b>Continue current order until insurance approved</b>

#### MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive rituximab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, newly diagnosed cardiac arrhythmias, severe abdominal pain or vomiting, and/or surgery.

#### PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

\*Per FDA labeling, Acetaminophen PO, Diphenhydramine IVP, and Methylprednisolone IVP is suggested prior to infusion

IV	Diphenhydramine	25mg	50mg		PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg		
	Famotidine	20mg	40mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
					Cetirizine	10mg				
					Loratadine	10mg				
					Other:					

#### SPECIFIC MEDICATION:

<input type="checkbox"/>	Rituxan	<b>Any rituximab biosimilar may be used according to payer guidelines</b>
<input type="checkbox"/>	Ruxience	
<input type="checkbox"/>	Truxima	
<input type="checkbox"/>	Riabni	

#### INDUCTION DOSE:

375mg/m<sup>2</sup> per 250 - 500ml NS IV to infuse per step protocol once weekly x 4 weeks

Other: \_\_\_\_\_

#### SPECIAL/LAB ORDERS:

\_\_\_\_\_

\_\_\_\_\_

#### MAINTENANCE DOSE: (begin \_\_\_\_\_ months after last induction dose)

500mg/500ml NS IV to infuse per step protocol

1000mg/500ml NS IV to infuse per step protocol

Other: \_\_\_\_\_

#### MAINTENANCE FREQUENCY:

Infuse dose every  4 months  6 months

Other: \_\_\_\_\_

Refills x 12 months unless noted otherwise here:

**Prescriber to monitor patient for symptoms of HBV infection and reactivation as clinically appropriate.**

#### LINE USE/CARE ORDERS:

Start PIV/Access CVC

Flush device per facility standard flushing procedure

Provide nursing care per Palmetto Infusion Nursing Procedures and post observation if indicated.

#### ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



#### PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

#### PRESCRIBER SIGNATURE: (No stamp signatures)

#### DATE

Dispense as written/Brand medically necessary	Substitution permitted