

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

Fabrazyme® (agalsidase beta) Standard Plan of Treatment

Rev 4.28.25

PATIENT DEMOGRAPHICS:

Patient Name:	
Patient's Phone:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG Gender: Allergies: See list NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

E75.21 - Fabry Disease
- Other:

REQUESTED DOCUMENTATION: PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	Serum IgG and GL-3 level	THERAPY:	
6	Current infusion rate (established patients)		
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: We may require a detailed Letter of Medical Necessity or clinical supporting documentation (depending on diagnosis), to be able to verify eligibility and payment for this treatment through Medicare and/or other insurance plans.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

*FDA labeling suggests pre-medication with an antihistamine and antipyretic in patients who experience infusion reactions.

IV	Diphenhydramine	25mg	50mg	Other:	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
					Cetirizine	10mg				
					Loratadine	10mg				
					Other:					

MEDICATION:

Fabrazyme® (agalsidase beta) IV in 50 - 500mL
After completion of infusion, flush line with 20ml of NS

Patients ≥ 30kg: Initial infusion should be administered at 15mg/hr. Increase in increments of 3-5mg/hr with subsequent infusions as tolerated, with a minimum infusion time of 1.5hr.
Patients weighing < 30kg: Infuse at a maximum rate of 15mg/hr.

SPECIAL/LAB ORDERS:

DOSE:

1mg/kg
 Other: _____

FREQUENCY:

Every 2 weeks
 Other: _____

Refills x 12 months unless noted otherwise here:

NURSING ORDERS: ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Start PIV/Access CVC
 Flush device per facility standard flushing procedure
 Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated.

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures) DATE:

Dispense as written/Brand medically necessary	Substitution Permitted	



Checklist for referrals to Palmetto Infusion: Fax referral to 1.866.872.8920

- Patient demographics - address, phone number, SS#, etc.**
- Insurance information - copy of the card(s) if possible**
- Plan of Treatment/Orders**
- Most recent physician office notes to include tried and failed therapies - all insurance companies that require a pre-authorization require the note. This includes Medicare/Medicaid HMOs.**
- Any lab results or other diagnostic procedures to support the diagnosis.**

Palmetto Infusion will complete insurance verification and submit all required clinical documentation to the patient's insurance company for eligibility.

Our office will notify you if any further information is required.

We will review financial responsibility with patient and refer them to any available co-pay assistance as required. Palmetto Infusion Call Center 800.809.1265. Thank you for your referral.

www.palmettoinfusion.com