

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

Simponi ARIA[®] (golimumab) Pediatric Standard Plan of Treatment

Rev 4.29.25

PATIENT DEMOGRAPHICS:

Patient Name:	
Patient's Phone:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG Gender: Allergies: See list NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

M08._____ - Juvenile idiopathic arthritis (JIA), polyarticular
L40.5_____ - Psoriatic Arthropathy
_____ - Other:

REQUESTED DOCUMENTATION:

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	REQUIRED: TB screening for new start	THERAPY:	
6	HBV screening/labs as required by payor		
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive golimumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new onset or deterioration neurological changes, and/or surgery

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV	Diphenhydramine	25mg	50mg	Other:	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
					Cetirizine	10mg				
					Loratadine	10mg				
					Other:					

MEDICATION/DOSE:

Simponi ARIA[®] (golimumab) 80 mg/m² diluted in 100 ml NS given IV to infuse over at least 30 minutes

FREQUENCY:

Induction: Given at 0 week and 4 weeks, and then every 8 weeks thereafter

Maintenance: Given every 8 weeks

Other: _____

SPECIAL/OTHER LAB ORDERS:

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment. Prescriber to monitor patient for symptoms of HBV and TB infection and reactivation as clinically appropriate.

Refills x 12 months unless noted otherwise here: _____

NURSING ORDERS:

Start PIV/Access CVC

Flush device per facility standard flushing procedure

Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

_____	_____
Dispense as written/Brand medically necessary	Substitution permitted