

Referral Status:	MRN:
New referral	Order change
Order Renewal	
Preferred clinic:	DAYS IN AIC: DAYS IN HOME:

Tzield® (teplizumab-mzvw) Plan of Treatment

PATIENT DEMOGRAPHICS:

Patient Name:	Patient's Phone:
Date of Birth:	Address:
Allergies:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
	See list
	NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

E10.9 - Type 1 diabetes mellitus without complications	-Other:
E10.8 - Type 1 diabetes mellitus with unspecified complications	

REQUESTED DOCUMENTATION:

REQUESTED DOCUMENTATION:	HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE? Yes or No
1 Insurance information	PLEASE STATE
2 H&P including tried and failed therapies	REQUIRED WASHOUT
3 Full medication list	FROM PREVIOUS
4 Required: Recent CBC with diff and LFTs	THERAPY:
	IF ORDER CHANGE:
	Continue current order until insurance approved

HOME SUPPLY ORDER: *Applicable to all home infusion patients subject to insurance approval

All supplies for vascular access line care, dressing kit, drug administration, adverse reaction kit, Infusion pump, IV pole, pole clamp etc. will be

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive teplizumab-mzvw if receiving antibiotics for active infectious process, active fever and/or suspected infection or surgery. If patient experiences severe cytokine release syndrome, consider temporarily pausing therapy for 1-2 days and review prescribing information before continuing therapy.

<input checked="" type="checkbox"/>	Home Anaphylaxis Kit: Dispense and administer for mild and severe reaction. Applicable to all home infusion patients.
2 - Epinephrine 1 mg/ml 1 ml	
2 - Diphenhydramine 50 mg/ml 2 ml vial	
Syringes, needles and 0.9% Normal Saline Flushes 10 mls to administer	
Complete Home Infusion Physician Standing Order for complete Home Infusion, Nursing, and Adverse Reaction Orders to be utilized in the event of an adverse reaction/anaphylaxis.	
Epinephrine administered IM per weight based dosing guide and Benadryl 25-50mg IVP to be administered by clinician in the home.	

*Per FDA labeling: Premedication of Acetaminophen or NSAID and antihistamine, and/or an antiemetic is suggested for at least the first 5 days of the 14-day treatment course

*Any selected premedication will only be given with infusions 1 through 5 unless otherwise designated here:

IV	Diphenhydramine	25mg	50mg	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg		Famotidine	20mg	40mg		
	Ondansetron	4mg	8mg		Diphenhydramine	25mg	50mg		
	Other:				Fexofenadine	60mg	180mg		
					Cetirizine	10mg			
					Loratadine	10mg			
					Other:				

DOSAGE:

Treatment Day	Dosage
Day 1	65 mcg/m2
Day 2	125 mcg/m2
Day 3	250 mcg/m2
Day 4	500 mcg/m2
Day 5-14	1030 mcg/m2

LAB ORDERS:

<input type="checkbox"/>	Draw CBC with diff & LFT on infusion days 5 and 8
<input type="checkbox"/>	Other: _____

MEDICATION:

Tzield® dose to be diluted in 25ml NS administered as an IV infusion over a minimum of 30-minutes. Administer once daily for 14 consecutive days.

Follow first 5 infusions with a 1-hour post infusion observation.

Physician responsible for all follow up lab monitoring.

Per FDA labeling, with AST/ALT > 5x ULN, bilirubin > 3x ULN, or Lymphocyte count <500/mcL lasting one week or longer, patient may be ineligible to receive Tzield®

SITE OF CARE:

<input type="checkbox"/> AIC	<input type="checkbox"/> AIC and Home with a nurse
------------------------------	----------------------------------------------------

*If site of care not indicated, PIS will coordinate with patient

SPECIAL/OTHER LAB ORDERS:

<input type="checkbox"/>	_____
<input type="checkbox"/>	_____

LINE USE/CARE ORDERS:

<input checked="" type="checkbox"/>	Start PIV/Access CVC
<input checked="" type="checkbox"/>	Flush PIV/Access per PIV/PICC/CVC protocol.
<input checked="" type="checkbox"/>	Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated.

Dispense and Administer as Prescribed

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted